arXiv:1210.2097v1 [math.NA] 7 Oct 2012

VARIATIONAL TIME DISCRETIZATION OF GEODESIC CALCULUS

MARTIN RUMPF* AND BENEDIKT WIRTH†

Abstract. We analyze a variational time discretization of geodesic calculus on finite- and certain classes of infinite-dimensional Riemannian manifolds. We investigate the fundamental properties of discrete geodesics, the associated discrete logarithm, discrete exponential maps, and discrete parallel transport, and we prove convergence to their continuous counterparts. The presented analysis is based on the direct methods in the calculus of variation, on Γ-convergence, and on weighted finite element error estimation. The convergence results of the discrete geodesic calculus are experimentally confirmed for a basic model on a two-dimensional Riemannian manifold.

This provides a theoretical basis for applications in computer vision: In particular, discrete geodesics offer an effective tool for shape morphing and the computation of a distance between shapes, the discrete logarithm allows a linear representation of strongly nonlinear shape variability, the discrete exponential map provides a robust tool for shape extrapolation, and the discrete parallel transport can be used to transfer geometric details from one shape to another. The basic operations are exemplarily illustrated for two different shape spaces, the space of viscous volumetric objects and the space of discrete viscous shells.

Key words. geodesics, exponential map, logarithm, parallel transport, finite elements, error analysis, shape space

AMS subject classifications. 37L65, 49M25, 53C22, 65L20, 65D18

1. Introduction.

Riemannian geometry is a powerful theory which allows to transfer many important concepts (shortest connecting paths, the arithmetic mean, or the principal component analysis, to name but a few examples) from linear vector spaces onto nonlinear, curved spaces. It is based on the notion of a Riemannian metric, and it provides a set of basic tools useful in applications, among which we shall concentrate on geodesics, geodesic distance, exponential and logarithmic maps, and parallel transport. During the past decade, Riemannian concepts have for instance increasingly been applied in computer vision for the design and investigation of nonlinear and often infinite-dimensional shape spaces, where the Riemannian metric encodes the preferred shape variability. Applications include shape morphing and modeling [17], computational anatomy in which the morphing path establishes correspondences between a patient and a template [3], as well as shape statistics [9].

Notwithstanding the conceptual power, operators such as the logarithm or the exponential map involve the solution of time-dependent nonlinear ordinary or partial differential equations. In more complex spaces—as they appear in vision applications—they are typically difficult to compute unless the spaces possess very peculiar structures [31, 29]. Therefore, we here develop a discrete theory that can be seen as a natural time-discretization of the above-mentioned Riemannian calculus and which is comparatively simple to state and to implement. It is centered around the definition of a discrete geodesic as the minimizer of a time-discrete path energy and naturally extends from there to discrete analogs of logarithm, exponential map, and parallel transport.

Complementarily to earlier work [26], which deals with one particular shape space and focuses on the experimental study of the resulting discrete calculus, we here provide the theoretical justification of the general approach in terms of a rigorous convergence analysis for decreasing time step size under suitable assumptions on the manifold and the functional involved in the approximation. Furthermore, we discuss the range of applications and the method's current limitations.

In what follows, let us give an overview on Riemannian calculus in the context of shape spaces in computer vision, which motivates our work. There is a rich diversity of Riemannian

---

[*] Bonn University, Endenicher Allee 60, D-53115 Bonn, Germany (`Martin.Rumpf@ins.uni-bonn.de`).

[†] Courant Institute, 251 Mercer Street, New York NY 10012, USA (`Benedikt.Wirth@cims.nyu.edu`).

shape spaces in the literature. While some of them are finite-dimensional and represent shapes as polygonal curves or triangulated surfaces [17, 20], most deal with infinite-dimensional shapes such as planar curves which require Riemannian metrics with regularizing properties, for instance curvature-based or Sobolev-type metrics [21, 28]. Interestingly, there is a strong link to mechanics. Based on the principle of least action, also complex mechanical systems like incompressible inviscid fluid flows can be rephrased in a Riemannian language, taking the kinetic energy as metric [2]. Exploiting this idea, the space of sufficiently smooth domains $\mathcal{O} \subset \mathbb{R}^d$ can be assigned a Riemannian metric by identifying the tangent space at $\mathcal{O}$ with velocity fields $v : \mathcal{O} \to \mathbb{R}^d$ and defining a (physical) metric on these. For instance, Dupuis *et al*. employ a metric $g(v, v) = \int_D Lv \cdot v \, \mathrm{d}x$ for a higher order elliptic operator L on some computational domain $D \subset \mathbb{R}^d$ [7], ensuring a diffeomorphism property of geodesic paths (*cf*. Section 3.3 below). In contrast, Fuchs *et al*. [11] and Wirth *et al*. [30] consider a second order elliptic operator based on viscous fluid flow, which is only defined on the shape domain $\mathcal{O}$. The proposed discrete geodesic calculus might also be of use in optimal transport, where for two given probability densities $\rho_0, \rho_1 : D \to \mathbb{R}$ the Monge–Kantorovich functional $\int_D |\psi(x) - x|^2 \rho_0(x) \, \mathrm{d}x$ is minimized over all mass preserving mappings $\psi : D \to D$. Via a flow reformulation due to Benamou and Brenier [5], this fits nicely into the Riemannian context.

There are only few nontrivial application-oriented Riemannian spaces in which geodesics can be computed in closed form (*e.g*. [31, 29]), else the system of geodesic ODEs has to be solved via time stepping (*e.g*. [18, 4]). Geodesics can also be obtained variationally by minimizing the discretized path length [27] or energy [11, 30], an approach generalized and simplified in this work.

The idea of variational time discretization, which underlies our discrete geodesic calculus, has proved very appropriate also in other fields, in particular in the discretization of gradient flows for an energy with respect to a particular (Riemannian) metric (*e.g*. mean curvature flow [1]). For Hamiltonian mechanical systems, variational time discretization is by now a classic field. The corresponding analog of our time-discrete path energy already occurred in the 70's as discrete action sum (*cf*. the brief historic account in [13]) and was analyzed as an independent, time-discrete system, whose discrete symplectic time steps exhibit the same structure as the discrete exponential proposed here. The consistency with the associated time-continuous mechanical systems was much later exploited numerically (*cf*. the general overview on variational, structure-preserving ODE integration [13]) and analyzed for particular systems from the Γ-convergence [22] and from the ODE-discretization perspective [19] under the name of variational integrators. These numerical schemes possess several advantages such as the (approximate) conservation of conserved physical quantities, the reproduction of the correct statistical properties, or the straightforward incorporation of additional system constraints, for which reason they have been applied in mechanical as well as computer vision contexts [12, 15]. More recently, time-adaptive versions have been developed [23], and the extremality of time-discrete action sums has also been applied as a constraint in optimal control [25].

Let us note that in contrast to the above works, we here lay emphasis on the Riemannian tools, in particular on computing a discrete geodesic between two fixed given points, on deriving a consistent discrete logarithm from this geodesic, and on using discrete exponential and logarithmic map to compute a consistent discrete parallel transport. In Section 2 we briefly set forth the different components of our discrete geodesic calculus after which we show its applications to exemplary shape spaces in Section 3. The existence and convergence properties of the discrete geodesics and corresponding operators are proved in Sections 5 and 6 after which we discuss how the theory applies on embedded manifolds and on certain shape spaces from Section 7.

2. Discrete geodesic calculus on a Riemannian manifold. In this section we present the concept of the discrete geodesic calculus, which comprises the notions of discrete geodesics, discrete logarithmic and exponential map, and discrete parallel transport. In preparation for this we briefly recall the corresponding continuous calculus. Precise assumptions on the manifold and involved approximating functionals will be given in the subsequent sections.

Let $(\mathcal{M}, g)$ be a smooth, complete Riemannian manifold, where g denotes the metric, given as a family of positive definite quadratic forms $g_y : T_y\mathcal{M} \times T_y\mathcal{M} \to \mathbb{R}$ for $y \in \mathcal{M}$, where $T_y\mathcal{M}$ denotes the tangent space in y. Given a smooth path $(y(t))_{t\in[0,1]}$ on $\mathcal{M}$, the length of this path is given by

$$\mathcal{L}[(y(t))_{t\in[0,1]}] = \int_0^1 \sqrt{g_{y(t)}(\dot{y}(t), \dot{y}(t))}\,\mathrm{d}t\,. \tag{2.1}$$

Given two points y_A and y_B in $\mathcal{M}$, the minimizer of $\mathcal{L}$ over all smooth paths $(y(t))_{t\in[0,1]}$ with $y(0) = y_A$ and $y(1) = y_B$ is a geodesic. More generally, geodesics are defined as local minimizers of the path length. Let us define the distance $\mathrm{dist}(y_A, y_B)$ between $y(0) = y_A$ and $y(1) = y_B$ as the minimal path length. After reparameterization, a length-minimizing geodesic is also a minimizer of the path energy

$$\mathcal{E}[(y(t))_{t\in[0,1]}] = \int_0^1 g_{y(t)}(\dot{y}(t), \dot{y}(t))\,\mathrm{d}t \tag{2.2}$$

and satisfies the constant speed property $g_{y(t)}(\dot{y}, \dot{y}) = \mathcal{L}^2[(y(t))_{t\in[0,1]}]$. The logarithm of a point y_B with respect to a point y_A is defined by $\log_{y_A} y_B = \dot{y}(0)$ if $(y(t))_{t\in[0,1]}$ is the unique shortest path with constant speed connecting $y(0) = y_A$ and $y(1) = y_B$. The exponential map $\exp_{y_A}$ maps every tangent vector $v \in T_{y_A}\mathcal{M}$ onto the endpoint of a geodesic starting at y_A with initial speed v, *i.e.* in the above notation, $\exp_{y_A} \dot{y}(0) = y(1) = y_B$. Furthermore, let us recall the parallel transport on a Riemannian manifold. Using the Levi-Civita connection ∇ a tangential vector field v is parallel along a curve $(y(t))_{t\in[0,1]}$ on $\mathcal{M}$ if $\nabla_{\dot{y}(t)} v(t) = 0$. If we sample a continuous path $(y(t))_{t\in[0,1]}$ at times $t_k = k\tau$ for $k = 0, \ldots, K$ and $\tau := \frac{1}{K}$, denoting $y_k = y(t_k)$, we obtain the following estimates for length and energy

$$\mathcal{L}[(y(t))_{t\in[0,1]}] \geq \sum_{k=1}^{K} \mathrm{dist}(y_{k-1}, y_k)\,, \qquad \mathcal{E}[(y(t))_{t\in[0,1]}] \geq \frac{1}{\tau} \sum_{k=1}^{K} \mathrm{dist}^2(y_{k-1}, y_k)\,, \tag{2.3}$$

where equality holds for geodesic paths due to the constant speed property. Indeed, the first estimate is straightforward, and application of Jensen's inequality yields the second estimate,

$$\sum_{k=1}^{K} \mathrm{dist}^2(y_{k-1}, y_k) \leq \sum_{k=1}^{K} \tau \int_{(k-1)\tau}^{k\tau} g_{y(t)}(\dot{y}(t), \dot{y}(t))\ \mathrm{d}t = \tau\, \mathcal{E}[(y(t))_{t\in[0,1]}]\,.$$

Based on these preliminaries we are now in the position to introduce a discrete geodesic calculus. The starting point is a local approximation of the squared Riemannian distance dist^2 by a functional $\mathcal{W}$. In detail, we suppose that a smooth functional $\mathcal{W} : \mathcal{M} \times \mathcal{M} \to \mathbb{R}$ is given such that

$$\mathrm{dist}^2(y, \tilde{y}) = \mathcal{W}[y, \tilde{y}] + O(\mathrm{dist}^3(y, \tilde{y})) \tag{2.4}$$

for all y, $\tilde{y} \in \mathcal{M}$. We will see later that $g_y = \frac{1}{2}\mathcal{W}_{,22}[y, y]$ implies (2.4) for smooth g and $\mathcal{W}$. Together with (2.3) this motivates the following definition of a *discrete path energy* and a *discrete path length* for discrete K-paths defined as $(K+1)$-tuples $(y_0, \ldots, y_K)$ with $y_k \in \mathcal{M}$ for $k = 0, \ldots, K$.

Definition 2.1 (Discrete length and energy). *For a discrete K-path $(y_0,\ldots,y_K)$ we define the* discrete length L *and the* discrete energy E *by*

$$\mathrm{L}[(y_0,\ldots,y_K)] = \sum_{k=1}^{K} \sqrt{\mathcal{W}[y_{k-1},y_k]}\,, \qquad \mathrm{E}[(y_0,\ldots,y_K)] = K\sum_{k=1}^{K} \mathcal{W}[y_{k-1},y_k]\,.$$

Then a discrete geodesic *(of order K) is defined as a minimizer of* $\mathrm{E}[(y_0,\ldots,y_K)]$ *for fixed end points y_0, y_K.*

To proceed with the definition of discrete logarithm and discrete exponential map we need to give a meaning to displacements $\tilde y - y$. Thus, in what follows let us assume that $\mathcal{M}$ is a subset of some Banach space $\mathbf{Y}$. This assumption is not restrictive because our constructions are purely local and can thus always be applied to charts of a more general manifold. Given a continuous constant speed geodesic $(y(t))_{t\in[0,1]}$ with $y(0) = y_A$ and $y(1) = y_B$ and a discrete geodesic $(y_0,\ldots,y_K)$ with $y_0 = y_A$ and $y_K = y_B$, we may view $y_1 - y_0$ as the discrete counterpart of $\tau\dot y(0)$ for $\tau = \frac{1}{K}$. Motivated by the fact that $\frac{1}{K}\log_{y_A}(y_B) = \tau\dot y(0)$ we hence give the following definition of a discrete logarithmic map.

Definition 2.2 (Discrete logarithm). *Suppose the discrete geodesic $(y_0,\ldots,y_K)$ is the unique minimizer of the discrete energy (2.1) with $y_0 = y_A$ and $y_K = y_B$, then we define the* discrete logarithm $\left(\frac{1}{K}\mathrm{LOG}\right)_{y_A}(y_B) = y_1 - y_0$. *Note that $\frac{1}{K}$ is part of the symbol and not a factor.*

In the special case $K = 1$ we obtain $\left(\frac{1}{K}\mathrm{LOG}\right)_{y_A}(y_B) = y_B - y_A$. As in the continuous case, the discrete logarithm can be considered as a *linear* representation (in the space $\mathbf{Y}$ of displacements on y_A) of the *nonlinear* variation y_B of y_A. As we will verify later under suitable assumptions, for a sequence of successively refined discrete geodesics we obtain $K\left(\frac{1}{K}\mathrm{LOG}\right)_y(\tilde y) \to \log_y(\tilde y)$ for $K\to\infty$.

In the continuous setting, the exponential map of a tangent vector $v \in T_{y_A}\mathcal{M}$ is defined via $\exp_{y_A} v = y_B$, where $y_B = y(1)$ for the (unique) geodesic $(y(t))_{t\in[0,1]}$ with $y(0) = y_A$ and $\dot y(0) = v$. Obviously, $\exp_{y_A}(\frac{k}{K}v) = y(\frac{k}{K})$ holds for $k = 0,\ldots,K$. We now aim at approximating $\exp_{y_A}(k\cdot)$ via a discrete counterpart $\mathrm{EXP}^k_{y_A}$ (the notation reflects the fact that on Lie-groups $\exp^k(\cdot) = \exp(k\cdot)$). The definition should be consistent with the discrete logarithm in the sense that $\mathrm{EXP}^k_{y_0}(\zeta_1) = y_k$ for a discrete geodesic $(y_0, y_1,\ldots,y_K)$ of order $K \geq k$ with $\zeta_1 = y_1 - y_0 = \left(\frac{1}{K}\mathrm{LOG}\right)_{y_0}(y_K)$. Our definition will reflect the following recursive properties of the continuous exponential map for $v \in T_y\mathcal{M}$ sufficiently small,

$$\begin{aligned}
&\exp_y(1v) = \left(\tfrac{1}{1}\log_y\right)^{-1}(v)\,, \quad \exp_y(2v) = \left(\tfrac{1}{2}\log_y\right)^{-1}(v)\,,\\
&\exp_y(kv) = \exp_{\exp_y((k-2)v)}(2v_{k-1}) \quad \text{for } v_{k-1} = \log_{\exp_y((k-2)v)}\exp_y((k-1)v)\,.
\end{aligned}$$

Replacing the tangent vector v by a displacement ζ, $\exp(k\cdot)$ by (EXP^k), and $\frac{1}{k}\log$ by $(\frac{1}{k}\mathrm{LOG})$ we obtain the following recursive definition of the discrete exponential map.

Definition 2.3 (Discrete exponential map). *For $\zeta \in \mathbf{Y}$ sufficiently small with $y+\zeta \in \mathcal{M}$ define*

$$\begin{aligned}
&\mathrm{EXP}^1_y(\zeta) = \left(\tfrac{1}{1}\mathrm{LOG}\right)_y^{-1}(\zeta)\,, \quad \mathrm{EXP}^2_y(\zeta) = \left(\tfrac{1}{2}\mathrm{LOG}\right)_y^{-1}(\zeta)\,,\\
&\mathrm{EXP}^k_y(\zeta) = \mathrm{EXP}^2_{\mathrm{EXP}^{k-2}_y(\zeta)}(\zeta_{k-1}) \quad \textit{with } \zeta_{k-1} = \left(\tfrac{1}{1}\mathrm{LOG}\right)_{\mathrm{EXP}^{k-2}_y(\zeta)}\mathrm{EXP}^{k-1}_y(\zeta)\,.
\end{aligned}$$

The smallness assumption on ζ ensures that $\left(\frac{1}{1}\mathrm{LOG}\right)_y$ and $\left(\frac{1}{2}\mathrm{LOG}\right)_y$ are invertible. It is straightforward to verify $\mathrm{EXP}^K_y = \left(\frac{1}{K}\mathrm{LOG}\right)_y^{-1}$ on the image of $\left(\frac{1}{K}\mathrm{LOG}\right)_y$. The central ingredient of this definition, the operator EXP_y^2 requires the solution of a problem with a variational constraint. Indeed, to compute $\mathrm{EXP}_y^2(\zeta)$, one considers all discrete geodesic paths (y, y_1, y_2) of order 2, where for any chosen y_2 the point y_1 must by definition be the unique minimizer of (2.1) so that we may write $y_1[y_2]$. Now, $\mathrm{EXP}_y^2(\zeta)$ is that point y_2 for which we have $y + \zeta = y_1[y_2]$, which can be equivalently described via the equation

$$y + \zeta = \mathrm{argmin}_{y_1 \in \mathcal{M}} \left(\mathcal{W}[y, y_1] + \mathcal{W}[y_1, y_2] \right) . \tag{2.5}$$

The corresponding Euler–Lagrange equation, as a nonlinear equation for $\mathrm{EXP}_y^2(\zeta) = y_2$, is given by $\mathcal{W}_{,2}[y, y + \zeta] + \mathcal{W}_{,1}[y + \zeta, y_2] = 0$.

REMARK 2.4 (Discrete geodesics versus discrete exponential shooting). *The variational definition* (2.5) *of the operator* EXP^2 *ensures that as long as a discrete geodesic connecting two end points* y_A, y_B *is unique, the discrete exponential map* $\mathrm{EXP}_{y_A}^k$ *applied to* $(\frac{1}{K}\mathrm{LOG})_{y_A}(y_B)$ *for* $k = 0, \ldots, K$ *retrieves exactly the same discrete geodesic as the minimization of the discrete energy* $\mathrm{E}[y_A, \hat{y}_1, \ldots, \hat{y}_{K-1}, y_B]$ *over all* $\hat{y}_1, \ldots, \hat{y}_{K-1}$, i.e. $y_k = \mathrm{EXP}_y^k(\frac{1}{K}\mathrm{LOG})_{y_A}(y_B)$, *where* $(y_A = x_0, y_1, \ldots, y_{K-1}, y_B = y_K)$ *is the discrete geodesic.*

Finally, we consider the discretization of parallel transport along a discrete (not necessarily geodesic) curve $(y_0, \ldots, y_K)$. In the continuous setting, given a curve $(y(t))_{t\in[0,1]}$ and a vector v_0 at $y(0)$, we define the parallel transport $\mathrm{P}_{(y(t))_{t\in[0,1]}} v_0$ of v_0 along the curve as the vector $v(1)$ resulting from the solution of the $\nabla_{\dot{y}(t)} v(t) = 0$ for $t \in [0,1]$ and initial data $v(0) = v_0$, where $\nabla_{\dot{y}}$ denotes the covariant derivative defined via the Levi-Civita connection ∇. There is a well-known first-order approximation of parallel transport called Schild's ladder [8, 16], which is based on the construction of a sequence of geodesic parallelograms. Given a curve $(y(t))_{t\in[0,1]}$ and a tangent vector $v_{k-1} \in T_{y((k-1)\tau)}\mathcal{M}$, the approximation $v_k \in T_{y(k\tau)}\mathcal{M}$ of the parallel transported vector via a geodesic parallelogram can be expressed as

$$y_{k-1}^p = \exp_{y((k-1)\tau)}(v_{k-1}) \,, \qquad y_k^c = \exp_{y_{k-1}^p}\left(\tfrac{1}{2}\log_{y_{k-1}^p}(y(k\tau))\right) ,$$
$$y_k^p = \exp_{y((k-1)\tau)}\left(2\log_{y((k-1)\tau)}(y_k^c)\right) , \qquad v_k = \log_{y(k\tau)}(y_k^p) \,.$$

Here, y_k^c is the midpoint of the two diagonals of the geodesic parallogramm with vertices $y((k-1)\tau)$, y_{k-1}^p, y_k^p, and $y(k\tau)$. This scheme can be easily transferred to discrete curves based on the discrete logarithm and the discrete exponential introduced before.

DEFINITION 2.5 (Discrete parallel transport). *Let* $(y_0, \ldots, y_K)$ *be a discrete curve in* $\mathcal{M}$ *with* $y_k - y_{k-1}$ *sufficiently small for* $k = 1, \ldots, K$*, then the* discrete parallel transport *of a displacement* ζ_0 *at* y_0 *along* $(y_0, \ldots, y_K)$ *is defined for* $k = 1, \ldots, K$ *via the iteration*

$$y_{k-1}^p = y_{k-1} + \zeta_{k-1} \,, \qquad y_k^c = y_{k-1}^p + \left((\tfrac{1}{2}\mathrm{LOG})_{y_{k-1}^p}(y_k)\right) ,$$
$$y_k^p = \mathrm{EXP}_{y_{k-1}}^2(y_k^c - y_{k-1}) \,, \qquad \zeta_k = y_k^p - y_k \,,$$

where ζ_k *is the transported displacement at* y_k*. We denote* $\mathbf{P}_{y_K,\ldots,y_0}\zeta_0 = \zeta_K$.

Above we have used $(\frac{1}{1}\mathrm{LOG})_y(\tilde{y}) = \tilde{y} - y$ as well as $\mathrm{EXP}_y^1(\zeta) = y + \zeta$. In the k^{th} step of the discrete parallel transport the Euler–Lagrange equations to determine y_k^c and ζ_k for given ζ_{k-1} and discrete curve $(y_0, \ldots, y_K)$ are

$$\mathcal{W}_{,2}[y_{k-1} + \zeta_{k-1}, y_k^c] + \mathcal{W}_{,1}[y_k^c, y_k] = 0 \,, \tag{2.6}$$
$$\mathcal{W}_{,2}[y_{k-1}, y_k^c] + \mathcal{W}_{,1}[y_k^c, y_k + \zeta_k] = 0 \,. \tag{2.7}$$

If $\mathcal{W}$ is symmetric, these conditions are the same as the Euler–Lagrange equations for backwards parallel transport so that $\mathbf{P}_{y_K,\ldots,y_0}^{-1} = \mathbf{P}_{y_0,\ldots,y_K}$, which however is generally not true if $\mathcal{W}$ is not symmetric. Note that for a continuous geodesic $y(t)$ the velocity $\dot{y}(t)$ at a time t equals the initial velocity $\dot{y}(0)$, parallel transported along the geodesic. In analogy, for a discrete geodesic $(y_0, \ldots, y_K)$ we have $y_{k+1} - y_k = \mathbf{P}_{y_k,\ldots,y_0}(y_1 - y_0)$ for $k = 0, \ldots, K-1$.

Given discrete parallel transport, the Levi-Civita connection $\nabla_\xi \eta$ for $\xi \in T_y\mathcal{M}$ and a vector field η in the tangent bundle $T\mathcal{M}$ can be also be approximated.

DEFINITION 2.6. *For $y \in \mathcal{M}$, small enough $\xi \in \mathbf{Y}$, and vectors η_0 attached to y and η_1 attached to $y + \xi$ (representing a discrete vector field), $\boldsymbol{\nabla}_\xi(\eta_0, \eta_1) = \mathbf{P}^{-1}_{y+\xi,y}\eta_1 - \eta_0$ defines a discrete connection.*

The inverse parallel transport satisfies the same Euler–Lagrange equations (2.6)-(2.7) as above (only this time y_k^c and ζ_{k-1} are determined from ζ_k). In analogy to $\nabla_{\dot{y}(t)}\dot{y}(t) = 0$ for a continuous geodesic a discrete geodesic $(y_0, \ldots, y_K)$ satisfies $\boldsymbol{\nabla}_{\Delta y_k}(\Delta y_k, \Delta y_{k+1}) = 0$ for $\Delta y_k = y_{k+1} - y_k$, $k = 0, \ldots, K-2$. If $\mathcal{W}$ is symmetric, we can also express the discrete connection as $\boldsymbol{\nabla}_\xi(\eta_0, \eta_1) = \mathbf{P}_{y,y+\xi}\eta_1 - \eta_0$.

3. Examples for shape space manifolds.

In this section we give some examples of Riemannian shape spaces used for shape analysis to motivate the discrete geodesic calculus investigated in this paper. Currently the full analysis in this paper applies to the finite-dimensional models in Sections 3.1 and 3.2 and the infinite-dimensional model in Section 3.2 as will be discussed in Section 7.

3.1. The space of viscous-fluid objects.

If we consider volumetric objects as shapes, *i.e.* open connected domains $\mathcal{O} \subset \mathbb{R}^d$, we are in general not interested in point to point correspondences between two different objects. Hence, if we restrict to objects $\mathcal{O}$ which are homeomorphic to a reference object $\hat{\mathcal{O}}$, then any shape $y = \mathcal{O}$ can be represented by an equivalence class of deformations $[\phi] = \{\tilde{\phi} \,|\, \tilde{\phi}(\hat{\mathcal{O}}) = \mathcal{O}\}$, and a path $(\mathcal{O}(t))_{t\in[0,1]}$ of shapes can be represented by multiple families $(\phi(t))_{t\in[0,1]}$ of deformations acting on the reference object $\hat{\mathcal{O}}$. A family $(\phi(t))_{t\in[0,1]}$ is associated with a (Eulerian) velocity field $v(t) = \dot{\phi}(t) \circ \phi^{-1}(t)$, and shape variations are equivalence classes of such motion fields with $\tilde{v} \sim v$ if $\tilde{v} \cdot n = v \cdot n$ on $\partial\mathcal{O}$, where n is the outer normal on $\partial\mathcal{O}$. One can define a physically motivated metric on shape variations as the minimal rate of dissipation in a Newtonian fluid occupying $\mathcal{O}$ when its free boundary moves according to the shape variation,

$$g_{\mathcal{O}}(v, v) = \min_{\{\tilde{v} \,|\, \tilde{v}\cdot n = v\cdot n \text{ on } \partial\mathcal{O}\}} \int_{\mathcal{O}} \tfrac{\lambda}{2}(\mathrm{tr}\epsilon[\tilde{v}])^2 + \mu\,\mathrm{tr}(\epsilon[\tilde{v}]^2)\,\mathrm{d}x\,, \tag{3.1}$$

where $\lambda, \mu > 0$, $\epsilon[v] = \frac{1}{2}(\nabla v + \nabla v^T)$. To approximate the squared distance between two objects $\mathcal{O}$ and $\tilde{\mathcal{O}}$ via a functional $\mathcal{W}$, we solve a minimization problem over all possible deformations between the two objects, *i.e.* $\mathcal{W}[\mathcal{O}, \tilde{\mathcal{O}}] = \min_{\psi(\mathcal{O})=\tilde{\mathcal{O}}} \int_{\mathcal{O}} W(A[\psi])\,\mathrm{d}x$ for some particular energy density W, where $A[\psi] = D\psi^T D\psi$ is the associated Cauchy–Green strain tensor. To ensure the requested consistency (2.4) of the functional $\mathcal{W}$ and the metric g we require $W(\mathbb{1}) = 0$, $DW(\mathbb{1}) = 0$, and $D^2W(\mathbb{1})(B, B) = \frac{\lambda}{4}(\mathrm{tr}B)^2 + \frac{\mu}{2}\mathrm{tr}(B^2)$ for all $B \in \mathbb{R}^{d,d}$ (*cf.* also Lemma 5.7 below). A suitable choice is $W(A) = \frac{\mu}{2}\mathrm{tr}A + \frac{\lambda}{4}\det A - \left(\mu + \frac{\lambda}{2}\right)\log\sqrt{\det A} - \frac{d\mu}{2} - \frac{\lambda}{4}$, which is rigid body motion invariant and isotropic so that $\mathcal{W}[Q\mathcal{O} + b, \tilde{Q}\tilde{\mathcal{O}} + \tilde{b}] = \mathcal{W}[\mathcal{O}, \tilde{\mathcal{O}}]$ for any two rotations and translations $Q, \tilde{Q} \in SO(d)$, $b, \tilde{b} \in \mathbb{R}^d$. Hence, the resulting discrete path energy is also invariant with respect to rigid body motions independent of the time step size τ (see [30, 26] for further details on this approach). Figures 3.1 and 3.2 show discrete geodesics between different shapes, where the colors indicate the local energy density W, which approximates the dissipation. The last three shapes in Figure 3.2 show extrapolations via the discrete exponential map. Figure 3.3 depicts an example of discrete parallel transport. Both in the metric $g_{\mathcal{O}}$ and in the functional



FIG. 3.1. *Discrete 8-geodesic between the digits 4 and 9 in the space of viscous fluid objects from Section 3.1. Colors indicate the local energy density W (from blue, low, to red, high), approximating viscous dissipation.*



FIG. 3.2. *Discrete geodesic between the digits 3 and 7, extrapolated beyond 7, for the same model as in Figure 3.1.*

$\mathcal{W}$ the boundary of the objects is not sufficiently controlled to ensure existence of optimal continuous or discrete paths. As a remedy in what follows we restrict the admissible shapes. Alternatively, one might add a suitable regularization for $\partial\mathcal{O}$ (*e.g.* a perimeter regularization, [30]) or combine the viscous-fluid approach for the object interior with the viscous shell approach (see Section 3.2) for its boundary.

Following Fuchs *et al.* [11] the previous model can be reduced to a finite-dimensional shape manifold by describing the boundary of $\mathcal{O}$ as a B-spline curve with n control points $q_1, \ldots, q_n$. Hence, each shape is represented by the vector $y = (q_1, \cdots, q_n)$. This time we pose the shortest path problem between two B-spline curves under the assumption of a point correspondence between these curves. The evaluation of the metric thus requires to solve the associated minimization problem (3.1) for given velocity of the control points $v = \dot{y} = (\dot{q}_1, \ldots, \dot{q}_n)$ (and thereby of all points on the curves). Instead of the nonlinear function $\mathcal{W}$ above we might consider for two sets of control points y and $\tilde{y}$ as in [11] the quadratic functional $\mathcal{W}[y, \tilde{y}] = \int_{\mathcal{O}[y]} \frac{\lambda}{2}(\mathrm{tr}\epsilon[\psi[y, \tilde{y}]])^2 + \mu \mathrm{tr}(\epsilon[\psi[y, \tilde{y}]]^2)\,\mathrm{d}x$ where $\mathcal{O}[y]$ is the object domain described by y and $\psi[y, \tilde{y}]$ is the energy-minimizing deformation under the Dirichlet boundary conditions given by the point to point correspondence between the two B-splines.

3.2. The space of viscous shells. Here we deal with shapes $\mathcal{S}$ which are shells of some small thickness $\delta > 0$. Deforming these thin material layers, the associated viscous friction is predominantly caused by two mechanisms, namely friction due to tangential (transversally uniform) in-plane deformation and friction due to (transversally non-uniform) deformation caused by bending. Assuming a point to point correspondence between two shells $\mathcal{S}$ and $\tilde{\mathcal{S}}$, similarly to before we can define their squared geodesic distance as the minimum energy dissipated during a distortion of $\mathcal{S}$ into $\tilde{\mathcal{S}}$. It can be approximated by a functional measuring the strain energy of a deformation $\phi : \mathcal{S} \to \mathbb{R}^d$ with $\phi(\mathcal{S}) = \tilde{\mathcal{S}}$. The tangential strain is described by the tangential Cauchy–Green strain tensor $A[\phi]$, which is defined pointwise as an endomorphism on the tangent space of $\mathcal{S}$ via $I(A[\phi]v, w) = \tilde{I}(d\phi(v), d\phi(w))$, where I and $\tilde{I}$ are the first fundamental forms on $\mathcal{S}$ and $\tilde{\mathcal{S}}$, respectively. As the corresponding energy we apply the same functional as in the volumetric cases above, this time acting on the tangential Cauchy–Green strain tensor, *i.e.* $\mathcal{W}_{\mathrm{tgl}}[y, \tilde{y}] = \delta \int_{\mathcal{S}} W(A[\phi])\,\mathrm{d}a$ for $\phi = \tilde{y} \circ y^{-1}$, where $y, \tilde{y}$ are parametrizations of $\mathcal{S}$ and $\tilde{\mathcal{S}}$, respectively, over a reference shell $\hat{\mathcal{S}}$. The dissipated energy due to bending can be approximated as follows. Let S, $\tilde{S}$ denote the shape operators on $\mathcal{S}$ and $\tilde{\mathcal{S}}$, respectively. Furthermore, define $S[\phi]$ as the pullback of the shape operator $\tilde{S}$ onto $\mathcal{S}$ given by $I(S[\phi]v, w) = \tilde{I}(\tilde{S}d\phi(v), d\phi(w))$. Now the bending energy of the deformation ϕ



FIG. 3.3. *The digit 4 is sheared or bulged, and the shearing and bulging variations are transported along the discrete geodesic to the digit 9 via discrete parallel transport based on the model from Section 3.1.*



FIG. 3.4. *Discrete 8-geodesic between two hand poses, based on the model of discrete viscous shells from Section 3.2. The colors indicate dissipation due to surface stretching (top) and bending (bottom).*

can be described via the difference $Q[\phi] = S[\phi] - S$, known as the *relative shape operator*, and it is expected to scale cubically in δ (*cf*. the rigorous limit analysis in [10]). For the sake of simplicity we choose $\mathcal{W}_{\text{bnd}}[y, \tilde{y}] = \delta^3 \int_{\mathcal{S}} |Q[\phi]|^2 \, \mathrm{d}a$, where $|Q| = \sqrt{\mathrm{tr} Q^T Q}$ denotes the Frobenius norm. Combining both energy contributions we finally obtain

$$\mathcal{W}[y, \tilde{y}] = \int_{\mathcal{S}} \delta W(A[\phi]) + \delta^3 |Q[\phi]|^2 \, \mathrm{d}a \tag{3.2}$$

with $\phi = \tilde{y} \circ y^{-1}$. To apply this approach in geometry processing we consider triangular meshes as discrete approximations of shells, where each shell is represented by a vector $y = (x_i)_{i=1,\ldots,n} \in \mathbb{R}^{nd}$ of n nodes $x_i \in \mathbb{R}^d$. The gradient $\nabla\phi$ of a deformation $\phi = \tilde{y} \circ y^{-1}$ is piecewise constant on each shell triangle so that $A[\phi]$ is also piecewise constant. To define a discrete shape operator on a triangle T we proceed as follows. For every edge $e = T_1 \cap T_2$ between two triangles, we obtain an edge normal N_e as the normalized sum of the unit normals on T_1 and T_2. Then, for a triangle T with cyclically ordered edges e_0, e_1, e_2 and corresponding edge normals N_0, N_1, N_2 we define the discrete shape operator as the piecewise constant endomorphism S on each triangle plane with $I(Se_k, e_k) = 2(N_{(k+1) \bmod 3} - N_{(k-1) \bmod 3})e_k$. From these discrete shape operators we then derive the piecewise constant relative shape operator as above. Finally, we obtain a discrete counterpart of the energy (3.2) as the basis of a time-discrete geodesic on the space of triangulated shells. For details on this approach we refer to [14]. Let us remark that this approach can be combined with the shape space model for viscous-fluid objects, where we simultaneously measure viscous dissipation on the object domain *and* its boundary contour. Figure 3.4 shows a discrete geodesic between two shells, color-coded with the local energy density in (3.2) due to both tangential and bending deformation.

If we restrict ourselves to one-dimensional shells in $\mathbb{R}^2$ and parametrize them over the interval $[0,1]$ we obtain

$$\mathcal{W}_{\mathrm{tgl}}[y,\tilde{y}] = \delta \int_0^1 W\left(\frac{|\tilde{y}_{,s}|^2}{|y_{,s}|^2}\right)|y_{,s}|\,\mathrm{d}s\,, \quad \mathcal{W}_{\mathrm{bnd}}[y,\tilde{y}] = \delta^3 \int_0^1 (\kappa[\tilde{y}]-\kappa[y])^2|y_{,s}|\,\mathrm{d}s$$

for the tangential distortion and the bending energy, where $\kappa[y] = \frac{1}{|y_{,s}|}\left(\frac{y_{,s}}{|y_{,s}|}\right)_{,s}\cdot\frac{D^{90}y_{,s}}{|y_{,s}|}$ is the curvature of a curve y, subscript $,s$ denotes the derivative, and D^{90} counterclockwise rotation by $\frac{\pi}{2}$, *i.e.* $n[y] = \frac{D^{90}y_{,s}}{|y_{,s}|}$ is the normal on the curve y. The underlying Riemannian metric g_y on the space of shell curves for which $\frac{1}{2}\mathcal{W}_{,22}[y,y] = g_y$ is given by

$$g_y(v,v) = \int_0^1 2\delta\frac{1}{|y_{,s}|}W''(1)|v_{,s}^{\mathrm{tgl}}|^2 + \delta^3\left(\partial_y\kappa[y](v)\right)^2|y_{,s}|\,\mathrm{d}s\,,$$

where $v_{,s}^{\mathrm{tgl}} = v_{,s}\cdot\frac{y_{,s}}{|y_{,s}|}$ denotes the tangential component of $v_{,s}$ and

$$\partial_y\kappa[y](v) = -\frac{\kappa[y]}{|y_{,s}|}v_{,s}^{\mathrm{tgl}} + \frac{1}{|y_{,s}|}\left(\frac{1}{|y_{,s}|}\left(\mathbb{1}-\frac{y_{,s}}{|y_{,s}|}\otimes\frac{y_{,s}}{|y_{,s}|}\right)v_{,s}\right)_{,s}\cdot n[y]\,.$$

Linearization of the bending energy and the choice $W(A) = \frac{(1-A)^2}{2}$ lead to a simplified model

$$\hat{\mathcal{W}}[y,\tilde{y}] = \int_0^1 \frac{\delta}{2}\left(1-\frac{|\tilde{y}_{,s}|^2}{|y_{,s}|^2}\right)^2|y_{,s}| + \delta^3(\tilde{y}_{,ss}-y_{,ss})^2|y_{,s}|\,\mathrm{d}s\,, \tag{3.3}$$

which corresponds via $g_y = \frac{1}{2}\mathcal{W}_{,22}[y,y]$ to the simplified metric

$$\hat{g}_y(v,v) = \int_0^1 2\delta\frac{|v_{,s}^{\mathrm{tgl}}|^2}{|y_{,s}|} + \delta^3|v_{,ss}|^2|y_{,s}|\,\mathrm{d}s \tag{3.4}$$

to which our convergence theory will apply as discussed in Section 7.

3.3. The flow of diffeomorphism approach. Here we consider a domain $\Omega\subset\mathbb{R}^d$ for $d=2,3$, and families of diffeomorphisms $(\phi(t))_{t\in[0,1]}:\bar{\Omega}\to\mathbb{R}^d$. In medical applications [3] the diffeomorphisms represent deformations of anatomic reference structures. Thus, each diffeomorphism $\phi:\bar{\Omega}\to\mathbb{R}^d$ represents a particular anatomic configuration or shape $y=\phi$ in the space $\mathcal{M}$ of all such diffeomorphisms. We associate the energy $\mathcal{E}[(\phi(t))_{t\in[0,1]}] = \int_0^1\int_\Omega Lv(t)\cdot v(t)\,\mathrm{d}x\,\mathrm{d}t$ with a path $(\phi(t))_{t\in[0,1]}$ in this shape space, where $v(t) = \dot{\phi}(t)\circ\phi(t)^{-1}$ represents the Eulerian velocity of the underlying flow and L is a higher order elliptic operator. Under suitable assumptions on L one can prove that paths of finite energy are indeed one-parameter families of diffeomorphisms as studied by Dupuis, Grenander and Miller [7]. Physically, the metric $g_{\phi(t)}(\dot{\phi}(t),\dot{\phi}(t)) = \int_\Omega Lv(t)\cdot v(t)\,\mathrm{d}x$ describes the viscous dissipation in a multipolar fluid model as investigated by Nečas and Šilhavý [24]. A specific model for the viscous dissipation is given by $g_\phi(\dot{\phi},\dot{\phi}) = \int_\Omega\frac{\lambda}{2}(\mathrm{tr}\epsilon[v])^2 + \mu\,\mathrm{tr}(\epsilon[v]^2) + \gamma|D^m v|^2\,\mathrm{d}x\,,$ $\lambda,\mu,\gamma>0$, $m>2$. The first two terms of the integrand again represent the usual dissipation density in a Newtonian fluid, whereas the third term represents a higher order measure for friction. Obviously, this model is invariant with respect to rigid body motions.

A suitable choice for the functional $\mathcal{W}$ approximating the associated squared distance is given by $\mathcal{W}[\phi,\tilde{\phi}] = \int_\Omega W(A[\psi])) + \gamma|D^m\psi|^2\,\mathrm{d}x$ for $\psi = \tilde{\phi}\circ\phi^{-1}$, where W is the energy density from Section 3.1. Once again, the resulting discrete path energy would be invariant with respect to rigid body motions.

4. Discrete geodesic calculus on embedded finite-dimensional manifolds. After having introduced a number of relatively complex shape manifolds as a motivation for its application, let us briefly demonstrate the discrete geodesic calculus for the much simpler case of an $(m-1)$-dimensional manifold $\mathcal{M}$ embedded in $\mathbb{R}^m$. We consider the simple energy $\mathcal{W}[y,\tilde{y}] = |\tilde{y}-y|^2$ which reflects the stored elastic energy in a spring spanned between points y and $\tilde{y}$ through the ambient space of $\mathcal{M}$ in $\mathbb{R}^m$. Thus, the discrete path energy is given by

$$\mathrm{E}[(y_0,\dots,y_K)] = K\sum_{k=1}^K |y_k - y_{k-1}|^2\,.$$

Now, we define the Lagrangian $F(X,\Lambda) = \mathrm{E}[(y_0,\dots,y_K)] - \Lambda\cdot D(X)$, where $X = (y_1,\dots,y_{K-1})$ are the actual point positions, $\Lambda = (\lambda_1,\dots,\lambda_{K-1})$ the Lagrange multipliers, and $D(X) = (d_{\mathcal{M}}(y_k))_{k=1,\dots,k-1}$ with $d_{\mathcal{M}}(\cdot)$ the (local) signed distance function of $\mathcal{M}$. Hence $\mathrm{grad}_X F(X,\Lambda) = 0$, $\mathrm{grad}_\Lambda F(X,\Lambda) = 0$ are the necessary conditions for $(y_0,\dots,y_K)$ to be a discrete geodesic with end points $y_0 = y_A$ and $y_K = y_B$. The resulting system of $(m+1)(K-1)$ degrees of freedom can be solved for instance by Newton's method. In particular, $(\mathrm{grad}_X F(X,\Lambda))_k = 2K\,(2y_k - y_{k-1} - y_{k+1}) - \lambda_k \nabla d_{\mathcal{M}}(y_k) = 0$ results in finding points y_k $(k=1,\dots,K-1)$ on $\mathcal{M}$ such that

$$2y_k - y_{k-1} - y_{k+1} \perp T_{y_k}\mathcal{M}\,, \tag{4.1}$$

while $\mathrm{grad}_\Lambda F(X,\Lambda) = D(X) = 0$ ensures that the points y_k are located on $\mathcal{M}$. Equation (4.1) is a discrete counterpart of the continuous geodesic equation $\ddot{y}(t) \perp T_{y(t)}\mathcal{M}$ for an arclength parametrized geodesic curve $(y(t))_{t\in\mathbb{R}}$ with $\frac{2y_k - y_{k-1} - y_{k+1}}{\tau^2} \approx -\ddot{y}(k\tau)$.

From these considerations, we derive for the discrete logarithm that $\zeta = (\frac{1}{2}\mathrm{LOG})_y\tilde{y}$ is described by the condition

$$\zeta - \tfrac{\tilde{y}-y}{2} \perp T_{y+\zeta}\mathcal{M}\,.$$

To achieve a simple geometric condition for the discrete exponential map $\mathrm{EXP}_y^2\zeta$ we consider a discrete geodesic (y,y_1,y_2) with $y_1 = y+\zeta$ and $y_2 = y_1+\eta$, where the displacement η is the actual degree of freedom. The necessary condition for (y,y_1,y_2) to be a discrete geodesic and thus for $y_2 = \mathrm{EXP}_y^2\zeta$ is

$$\zeta - \eta \perp T_{y+\zeta}\mathcal{M}\,,$$

which results in a one-dimensional search problem in the space $\mathcal{M}\cap(y+\mathrm{span}\{\zeta, n_{y+\zeta}\})$, where $n_{y+\zeta}$ denotes a normal on $T_{y+\zeta}\mathcal{M}$.

5. Properties of the discrete path energy. In the remainder of the paper we aim to examine the convergence properties of the discrete geodesic calculus as the discrete time step size $\tau = \frac{1}{K}$ tends to 0. Figure 5.1 shows experimental evidence for the convergence of the discrete geodesic, exponential map, logarithmic map, and parallel transport. In this section we study some properties of the discrete and the continuous path energy on a specific class of manifolds, and we examine their relation via the concept of Γ-convergence. In particular we will show that sequences of successively refined discrete geodesic paths converge to a continuous geodesic path.

We consider the following functional analytic set up. Let $\mathbf{Y}$ be a Banach space with norm $\|\cdot\|_{\mathbf{Y}}$ on which a symmetric bilinear form $(y,v,w)\mapsto g_y(v,w)$ for $y,\ v,\ w\in\mathbf{Y}$ induces a Riemannian structure with $g:\mathbf{Y}\times\mathbf{Y}\times\mathbf{Y}\to\mathbb{R}\cup\{\infty\}$. Furthermore, let $\mathbf{V}$ be a compactly embedded, separable, reflexive subspace of $\mathbf{Y}$ with norm $\|\cdot\|_{\mathbf{V}}$. We shall suppose that g is uniformly bounded and $\mathbf{V}$-coercive in the sense

$$c^*\|v\|_{\mathbf{V}}^2 \le g_y(v,v) \le C^*\|v\|_{\mathbf{V}}^2 \tag{5.1}$$



FIG. 5.1. *Top: Discrete geodesic path* $y_\tau = (y_0, \ldots, y_K)$ *with* $y_0 = y_A = (\frac{1}{2}, 0)$ *and* $y_1 = y_B = (-\frac{1}{2}, 2)$ *on the stereographic projection of the sphere,* $K(\frac{1}{K}\mathrm{LOG})_{y_A}(y_B)$*,* $(\mathrm{EXP}^K)_{y_A}(v/K)$ *for* $v = \log_{y_A}(y_B)$*, and parallel transport of* $w = (-\frac{2}{5}, 0)$ *from* y_A *to* y_B *at different temporal resolutions (*$K = 2^k$ *for* $k = 1, \ldots, 10$*). Bottom: The corresponding errors versus* $1/K$*.*

for constants $c^* > 0$ and $C^* < \infty$ independent of y. In particular, $g_y(v, v) = \infty$ if $v \notin \mathbf{V}$. Also, g shall be continuous in y with

$$|g_y(v, v) - g_{\tilde{y}}(v, v)| \leq \beta(\|y - \tilde{y}\|_{\mathbf{Y}})\|v\|_{\mathbf{V}}^2 \tag{5.2}$$

for a strictly increasing, continuous function β with $\beta(0) = 0$. Finally, given $y_A \in \mathbf{Y}$ we consider the manifold $\mathcal{M} := y_A + \mathbf{V}$, on which $g_\cdot(\cdot, \cdot)$ induces a Riemannian structure.

REMARK 5.1. *As a motivation for this setup, consider the infinite-dimensional shape space of 1D shells discussed in Section 3.2 (the application to finite-dimensional shape spaces is explained in the next remark). Shells* y *can be considered as differentiable curves in the space* $\mathbf{Y} = C^1([0, 1]; \mathbb{R}^2)$*, and the space* $\mathbf{V} = W^{2,2}((0, 1); \mathbb{R}^2)$ *encodes the regularity of admissible variations* v *of a shell* y *for which the metric* $g_y(v, v)$ *is finite. The shell manifold* $\mathcal{M}$ *is locally spanned by all those shapes which can be reached from a particular* $y_A \in \mathbf{Y}$ *via (short time) integration of* $\dot{y}(t) = v(t)$ *with* $v(t) \in W^{2,2}((0, 1); \mathbb{R}^2)$*.*

REMARK 5.2. *The assumptions particularly cover the local theory of smooth* k*-dimensional shape manifolds (*$k < \infty$*), which by Nash's theorem can be isometrically embedded in the Euclidian space* $\mathbb{R}^m$ *for* m *sufficiently large. In this case one chooses* $\mathbf{Y} = \mathbf{V} = \mathbb{R}^k$ *to represent a chart of the manifold and* $g_y(v, v) = DX(y)v \cdot DX(y)v$*, where* X *is the associated parametrization and* $\cdot$ *is the scalar product in* $\mathbb{R}^m$*.*

For y_A, $y_B \in \mathcal{M}$, the next theorem states the existence of a connecting path with least energy. The key point is the weak lower semi-continuity of the continuous path energy (2.2) using the compact embedding of $\mathbf{V}$ into $\mathbf{Y}$.

THEOREM 5.3 (Existence of continuous geodesics). *Under the above assumptions there exists a continuous geodesic path* $(y(t))_{t\in[0,1]}$ *defined as a minimizer of the path energy* $\mathcal{E}$ *among all paths* $(\tilde{y}(t))_{t\in[0,1]}$ *with* $\tilde{y}(0) = y_A$ *and* $\tilde{y}(1) = y_B$*.*

Proof. Let $(y^j(t))_{t\in[0,1]}$ be a minimizing sequence of paths. We can obviously assume that the energy on this sequence is bounded by $\bar{\mathcal{E}} = \int_0^1 g_{y_A + t(y_B - y_A)}(y_B - y_A, y_B - y_A)\,\mathrm{d}t$. Hence, $\|v^j\|_{L^2((0,1);\mathbf{V})}$ is bounded for $v^j = \dot{y}^j$. Because $L^2((0, 1); \mathbf{V})$ is separable and reflexive, there is a subsequence—with a slight misuse of notation indexed as before—which converges weakly to some $v \in L^2((0,1);\mathbf{V})$. Likewise, the curves $y^j(t) = \int_0^t v^j(s)\,\mathrm{d}s$ converge weakly in $W^{1,2}((0,1);y_A+\mathbf{V})$ to $y(t) = \int_0^t v(s)\,\mathrm{d}s$. Due to the bounded embedding of $W^{1,2}((0,1);\mathbf{V})$ into $C^{0,\frac{1}{2}}([0,1];\mathbf{V})$ and the compact embedding of $C^{0,\frac{1}{2}}([0,1];\mathbf{V})$ into $C^0([0,1];\mathbf{Y})$ by Arzelà–Ascoli, upon selecting a further subsequence, y^j converges strongly in $C^0([0,1];\mathbf{Y})$. Now choose j large enough that $\mathcal{E}[y^j] \leq \underline{\mathcal{E}} + \delta$ and $\sup_{t\in[0,1]} \beta(\|y^j(t) - y(t)\|_\mathbf{Y}) \leq \delta$ for a prescribed small $\delta > 0$ and $\underline{\mathcal{E}} = \inf \mathcal{E}$. Following the usual paradigm of the direct methods in the calculus of variations we can estimate the energy using Mazur's lemma, the convexity of $v \mapsto g_y(v,v)$, the continuity and coercivity of g, and Fatou's lemma,

$$\begin{aligned}
\mathcal{E}[(y(t))_{t\in[0,1]}] &= \int_0^1 g_{y(t)}(v(t),v(t))\,\mathrm{d}t \\
&\leq \int_0^1 \lim_{j\to\infty} \sum_{i=j}^{N_j} \lambda_i^j g_{y(t)}(v^i(t),v^i(t))\,\mathrm{d}t \\
&\leq \int_0^1 \lim_{j\to\infty} \sum_{i=j}^{N_j} \lambda_i^j g_{y^i(t)}(v^i(t),v^i(t)) + \lambda_i^j \beta(\|y^i(t)-y(t)\|_\mathbf{Y})\|v^i(t)\|_\mathbf{V}^2\,\mathrm{d}t \\
&\leq \lim_{j\to\infty} \sum_{i=j}^{N_j} \lambda_i^j \left(\mathcal{E}[(y^i(t))_{t\in[0,1]}] + \delta \int_0^1 \|v^i(t)\|_\mathbf{V}^2\,\mathrm{d}t \right) \leq \underline{\mathcal{E}}\left(1 + \frac{\delta}{c^*}\right).
\end{aligned}$$

Here, $(N_j)_{j=1,\ldots,\infty}$ is a sequence in $\mathbb{N}$ and $(\lambda_i^j)_{i=j,\ldots,N_j}^{j=1,\ldots,\infty}$ is a sequence of convex combination coefficients with $\sum_{i=j}^{N_j} \lambda_i^j = 1$, $\lambda_i^j \geq 0$, and $\sum_{i=j}^{N_j} \lambda_i^j v^i \to v$ strongly in $L^2((0,1);\mathbf{V})$ for $j \to \infty$. Because δ is arbitrary, we obtain $\mathcal{E}[(y(t))_{t\in[0,1]}] \leq \underline{\mathcal{E}}$, which proves the claim. □

For y_A, $y_B \in \mathbf{Y}$, let us introduce the Riemannian distance

$$\operatorname{dist}(y_A,y_B) = \sqrt{\min_{y(0)=y_A,y(1)=y_B} \mathcal{E}[(y(t))_{t\in[0,1]}]}\,. \tag{5.3}$$

It is an easy exercise to verify the axioms of a metric and that the induced topology is equivalent to the $\mathbf{V}$-topology, $\sqrt{c^*}\|y_B - y_A\|_\mathbf{V} \leq \operatorname{dist}(y_A,y_B) \leq \sqrt{C^*}\|y_B - y_A\|_\mathbf{V}$. Furthermore, a simple reparameterization argument shows $g_{y(t)}(\dot{y}(t),\dot{y}(t)) = \text{const.}$ along a geodesic.

For the discrete path energy, we would like to show an analogous existence result, as well as properties related to (5.3) and the above-mentioned constant speed parameterization. For this purpose we consider a lower semi-continuous $\mathcal{W} : \mathbf{Y} \times \mathbf{Y} \to \mathbb{R} \cup \{\infty\}$ satisfying

$$\mathcal{W}[y,\tilde{y}] = \operatorname{dist}^2(y,\tilde{y}) + O(\operatorname{dist}^3(y,\tilde{y})) \tag{5.4}$$

with uniform constants, and we assume coercivity of $\mathcal{W}$ in the sense

$$\mathcal{W}[y,\tilde{y}] \geq \gamma(\operatorname{dist}(y,\tilde{y})) \tag{5.5}$$

for a strictly increasing, continuous function γ with $\gamma(0) = 0$.

THEOREM 5.4 (Existence of discrete geodesics). *Given y_A, $y_B \in \mathcal{M}$, there is a discrete geodesic path $(y_0,\ldots,y_K)$ which minimizes the discrete energy E over all discrete paths $(\tilde{y}_0,\ldots,\tilde{y}_K)$ with $\tilde{y}_0 = y_A$ and $\tilde{y}_K = y_B$.*

Proof. Let $((y_0^j,\ldots,y_K^j))_{j=1,\ldots,\infty}$ be a minimizing sequence. We can obviously assume the y_k^j to lie on $\mathcal{M} = y_A + \mathbf{V}$ and that the energy on this sequence is bounded by $\bar{\mathrm{E}} = K\mathcal{W}[y_A,y_B] + K(K-1)\mathcal{W}[y_B,y_B]$. Because $\mathbf{V}$ compactly embeds in $\mathbf{Y}$, there is a subsequence, still denoted $(y_0^j,\ldots,y_K^j)$, which converges in $\mathbf{Y}$ to a discrete path $(y_0,\ldots,y_K)$.

By the lower semi-continuity of $\mathcal{W}$ in both arguments, we finally obtain $\mathrm{E}[(y_0,\dots,y_K)] \leq \inf_{(\tilde y_0,\dots,\tilde y_K)} \mathrm{E}[(\tilde y_0,\dots,\tilde y_K)]$. □

THEOREM 5.5 (Bounds on discrete path energy). *The discrete path energy satisfies*

$$\min_{\substack{(y_0,\dots,y_K)\\ y_0=y_A, y_K=y_B}} \mathrm{E}[(y_0,\dots,y_K)] = \mathrm{dist}^2(y_A,y_B)(1+O(\mathrm{dist}(y_A,y_B)/K))$$

with uniform constants.

Proof. Denote the minimum energy value by $\underline{\mathrm{E}}$. Choosing points $y_1,\dots,y_{K-1}$ such that $\mathrm{dist}(y_{k-1},y_k) = \mathrm{dist}(y_0,y_K)/K$ for $k=1,\dots,K$, we directly see

$$\underline{\mathrm{E}} \leq K \textstyle\sum_{k=1}^K \mathcal{W}[y_{k-1},y_k] = \mathrm{dist}^2(y_0,y_K)(1+O(\mathrm{dist}(y_0,y_K)/K))\,.$$

On the other hand, letting $a_k = \mathrm{dist}(y_{k-1},y_k)$ for the minimizing $(y_0,\dots,y_K)$, we have $\underline{\mathrm{E}} \geq K\sum_{k=1}^K a_k^2 - Ca_k^3 =: F(a_1,\dots,a_K)$ for some $C>0$. From the coercivity of $\mathcal{W}$ we know $\underline{\mathrm{E}} \geq K\sum_{k=1}^K \gamma(a_k)$ and thus $a_k \leq \gamma^{-1}(\mathrm{dist}^2(y_0,y_K)(\frac{1}{K}+O(\frac{\mathrm{dist}(y_0,y_K)}{K^2})))$. Thus, for K large enough, $a_k \leq 1/3C$ for all k. Minimization of F under the constraints $\sum_{k=1}^K a_k \geq \mathrm{dist}(y_0,y_K)$ and $a_k \leq 1/3C$, $k=1,\dots,K$, yields $a_1 = \ldots = a_K = \mathrm{dist}(y_0,y_K)/K$ and thus $\underline{\mathrm{E}} \geq F(a_1,\dots,a_K) = \mathrm{dist}^2(y_0,y_K)(1-C\mathrm{dist}(y_0,y_K)/K)$. □

THEOREM 5.6 (Equidistribution of points along discrete geodesics). *Discrete geodesics (minimizers from Theorem 5.4) satisfy* $\mathrm{dist}(y_{k-1},y_k) \leq C\mathrm{dist}(y_A,y_B)/K$ *for* $k=1,\dots,K$.

Proof. For given K, let $j_K \in \{1,\dots,K\}$ be such that $\gamma_K := \mathrm{dist}(y_{j_K-1},y_{j_K})$ is largest. Furthermore abbreviate $d := \mathrm{dist}(y_0,y_K)$ and $\alpha_K := \mathrm{dist}(y_0,y_{j_K-1})/d$. We have

$$\begin{aligned}
&d^2(1+O(d/K)) = \mathrm{E}[(y_0,\dots,y_K)]\\
&\quad\geq \tfrac{K}{j_K-1}\mathrm{E}[(y_0,\dots,y_{j_K-1})] + K\gamma_K^2(1-O(\gamma_K)) + \tfrac{K}{K-j_K}\mathrm{E}[(y_{j_K},\dots,y_K)]\\
&\quad\geq K\gamma_K^2(1-O(\gamma_K)) + \tfrac{K}{j_K-1}\mathrm{dist}^2(y_0,y_{j_K-1})(1-O(\gamma_K)) + \tfrac{K}{K-j_K}\mathrm{dist}^2(y_{j_K},y_K)(1-O(\gamma_K))\\
&\quad\geq (1-O(\gamma_K))(K\gamma_K^2 + d^2[\tfrac{K}{j_K-1}\alpha_K^2 + \tfrac{K}{K-j_K}(1-\alpha_K-\tfrac{\gamma_K}{d})^2])\,.
\end{aligned}$$

This is minimized by $\alpha_K = \frac{(j_K-1)(1-\frac{\gamma_K}{d})}{K-1}$ and yields

$$d^2(1+O(d/K)) \geq (1-O(\gamma_K))(K\gamma_K^2 + d^2\tfrac{K}{K-1}(1-\tfrac{\gamma_K}{d})^2)\,.$$

For $\gamma_K = \frac{\omega_K}{K}$ with $\omega_K \to \infty$ for $K \to \infty$ and $\omega_K \leq C\sqrt{K}$ by the trivial estimate $\gamma_K \leq \frac{C}{\sqrt{K}}$ one obtains $d^2 + \frac{Cd^3}{K} \geq (1-\frac{C}{\sqrt{K}})(\frac{\omega_K^2}{K} + d^2\frac{K}{K-1}(1-\frac{\omega_K}{dK})^2)$ which yields a contradiction, so $\mathrm{dist}(y_{j_K-1},y_{j_K}) \leq Cd/K$. □

For the subsequent estimates it is convenient to relate the function $\mathcal{W}$ to the metric g; we use the following.

LEMMA 5.7 (Consistency conditions). *If* $v \mapsto \mathcal{W}[y,y+v]$ *is twice Gâteaux-differentiable on* $\mathbf{V}$ *for* $y \in \mathbf{Y}$*, then* $\mathcal{W}[y,\tilde y] = \mathrm{dist}^2(y,\tilde y) + O(\mathrm{dist}^3(y,\tilde y))$ *for* $\tilde y$ *close to* y *implies*

$$\mathcal{W}[y,y] = 0\,,\quad \mathcal{W}_{,2}[y,y](v) = 0\,,\quad \mathcal{W}_{,22}[y,y](v,w) = 2g_y(v,w)$$

for any $v,w \in \mathbf{V}$*. Furthermore,* $\mathcal{W}_{,1}[y,y](v) = 0$ *and*

$$\mathcal{W}_{,11}[y,y](v,w) = -\mathcal{W}_{,12}[y,y](v,w) = -\mathcal{W}_{,21}[y,y](v,w) = \mathcal{W}_{,22}[y,y](v,w)\,.$$

If $v \mapsto \mathcal{W}[y,y+v]$ *is even three times Fréchet-differentiable, the implication becomes an equivalence.*

Proof. It is readily shown that $\mathrm{dist}^2(y,\tilde y) = g_y(\tilde y - y, \tilde y - y) + O(\beta(\|\tilde y - y\|_{\mathbf{Y}}))\|\tilde y - y\|_{\mathbf{V}}^2$.

Let $\mathcal{W}[y,\tilde y] = \mathrm{dist}^2(y,\tilde y) + O(\mathrm{dist}^3(y,\tilde y))$ for $\tilde y$ close to y, then obviously $\mathcal{W}[y,y] = 0$ and $\mathcal{W}_{,2}[y,y](v) = \frac{\mathrm{d}}{\mathrm{d}t}\mathcal{W}[y,y+tv] = 0$. Now assume $\mathcal{W}_{,22}[y,y](v,v) \neq 2g_y(v,v)$ for some $v \in \mathbf{V}$ (note that due to the bilinearity it is sufficient to show equality of the quadratic forms). Without loss of generality, let $\mathcal{W}_{,22}[y,y](v,v) > \alpha g_y(v,v)$ for some $\alpha > 2$. This implies $\mathcal{W}[y,y+tv] > \frac{1}{2}\left(\frac{\alpha}{2}+1\right) g_y(tv,tv)$ for t small enough, which (using $\mathrm{dist}^2(y,y+tv) = g_y(tv,tv) + o(t^2)$) is strictly greater than $(\frac{\alpha}{4}+\frac{1}{2})\mathrm{dist}^2(y,y+tv) + o(\mathrm{dist}^2(y,y+tv))$ and thus a contradiction.

The above applied to the first argument of $\mathcal{W}$ instead of the second implies $\mathcal{W}_{,1}[y,y] = 0$ and $\mathcal{W}_{,11}[y,y](v,w) = 2g_y(v,w)$. Finally, for any curve $(y(t))_{t\in\mathbb{R}}$ in $\mathbf{Y}$ we can differentiate $0 = \mathcal{W}_{,1}[y(t),y(t)]$ and $0 = \mathcal{W}_{,2}[y(t),y(t)]$ with respect to t, yielding $\mathcal{W}_{,12}[y,y] = -\mathcal{W}_{,11}[y,y]$ and $\mathcal{W}_{,21}[y,y] = -\mathcal{W}_{,22}[y,y]$.

If $v \mapsto \mathcal{W}[y,y+v]$ is three times Fréchet-differentiable and $\mathcal{W}_{,22}[y,y](v,w) = 2g_y(v,w)$, then by Taylor's theorem for $\tilde y = y + tv$, $\mathcal{W}[y,\tilde y] = g_y(tv,tv) + O(t^3) = \mathrm{dist}^2(y,\tilde y) + O(\mathrm{dist}^3(y,\tilde y))$. □

THEOREM 5.8 (Uniqueness of discrete geodesics). *If $v \mapsto \mathcal{W}[y,y+v]$ is twice Gâteaux-differentiable on $\mathbf{V}$ for $y \in \mathcal{M}$ and if $\|y_B - y_A\|_{\mathbf{V}}$ is sufficiently small there exists a unique discrete geodesic $(y_0,\ldots,y_K)$ with $y_0 = y_A$ and $y_K = y_B$.*

Proof. Without any restriction we assume $y_A \in \mathbf{V}$. Otherwise, we have to work with offsets $y_k - y_A$ instead of points y_k. For $X_{0,K} := (y_0, y_K)$ and $X_{1,\ldots,K-1} := (y_1,\ldots,y_{K-1})$ we define the function $F : \mathcal{M}^{K+1} \to (\mathbf{V}')^{K-1}; (X_{0,K}, X_{1,\ldots,K-1}) \mapsto (\mathcal{W}_{,2}[y_{k-1},y_k] + \mathcal{W}_{,1}[y_k,y_{k+1}])_{k=1,\ldots,K-1}$, where $\mathbf{V}'$ denotes the dual space of $\mathbf{V}$. Then the block tridiagonal operator $\mathcal{A} = D_{X_{1,\ldots,K-1}}F(X_{0,K},X_{1,\ldots,K-1})$ is a $(K-1)\times(K-1)$ is given by

$$\begin{aligned}\mathcal{A}_{kk} &= \mathcal{W}_{,22}[y_{k-1},y_k] + \mathcal{W}_{,11}[y_k,y_{k+1}]\,,\\ \mathcal{A}_{k(k-1)} &= \mathcal{W}_{,21}[y_{k-1},y_k]\,, \quad \mathcal{A}_{k(k+1)} = \mathcal{W}_{,12}[y_k,y_{k+1}]\,.\end{aligned}$$

Now, using Lemma 5.7 we obtain for the trivial geodesic $(y_A,\ldots,y_A)$ that

$$\mathcal{A} = \begin{pmatrix} 2g & -g & & & \\ -g & 2g & -g & & \\ & \ddots & \ddots & \ddots & \\ & & -g & 2g & -g \\ & & & -g & 2g \end{pmatrix} = \mathrm{diag}(g^{\frac{1}{2}}) \begin{pmatrix} 2\mathbb{1} & -\mathbb{1} & & & \\ -\mathbb{1} & 2\mathbb{1} & -\mathbb{1} & & \\ & \ddots & \ddots & \ddots & \\ & & -\mathbb{1} & 2\mathbb{1} & -\mathbb{1} \\ & & & -\mathbb{1} & 2\mathbb{1} \end{pmatrix} \mathrm{diag}(g^{\frac{1}{2}})\,,$$

where $g = g_{y_A}$. Hence, the inverse of $\mathcal{A}$ can be computed by Gaussian elimination, which ensures that $\mathcal{A} = D_{X_{1,\ldots,K-1}}F(y_A,\ldots,y_A)$ is invertible with bounded inverse, and thus the claim follows by the implicit function theorem. □

In what follows, we aim to prove convergence of discrete geodesics against continuous ones. To this end we identify any discrete path $(y_0,\ldots,y_K)$ on $\mathcal{M}$ with its piecewise geodesic interpolation $\hat y^K : [0,1] \to \mathcal{M}$, *i.e.* every segment $\hat y^K|_{[k\tau,(k+1)\tau]}$ shall be the shortest continuous connecting geodesic between y_k and y_{k+1}. Now, we define an energy E^K on continuous paths via $\mathrm{E}^K[(y(t))_{t\in[0,1]}] := \mathrm{E}[(y_0,\ldots,y_K)]$ if $y(t) = \hat y^K(t)$ for some $y_0,\ldots,y_K \in \mathcal{M}$ with $y_0 = y_A$, $y_K = y_B$, and $\mathrm{E}^K[(y(t))_{t\in[0,1]}] = \infty$ else. Based on these notational preliminaries we obtain the following convergence result.

THEOREM 5.9 (Γ-convergence of the discrete energy). *In the $L^2((0,1);\mathbf{Y})$-topology the Γ-limit of E^K for $K \to \infty$ is $\mathcal{E}$.*

Proof. To verify Γ-convergence we have to establish the two defining properties, the limsup- and the liminf-inequality.

[*limsup-inequality*] For an arbitrary $y : [0,1] \to \mathcal{M}$ with $y \in L^2((0,1);\mathbf{Y})$ we have to show that there exists a sequence of curves $\hat y^K : [0,1] \to \mathcal{M}$ with $\hat y^K \to y$ in $L^2((0,1);\mathbf{Y})$ and

$\limsup_{K\to\infty} \mathrm{E}^K[(\hat y^K(t))_{t\in[0,1]}] \leq \mathcal{E}[(y(t))_{t\in[0,1]}]$. Without any restriction we can assume $\mathcal{E}[(y(t))_{t\in[0,1]}] < \infty$. Applying the Cauchy–Schwarz inequality we get

$$\mathrm{dist}(y(t),y(s)) = \int_t^s \sqrt{g_{y(r)}(\dot y(r),\dot y(r))}\,\mathrm{d}r \leq \sqrt{|s-t| \int_t^s g_{y(r)}(\dot y(r),\dot y(r))\,\mathrm{d}r}\,, \tag{5.6}$$

which immediately implies $W^{1,2}((0,1);\mathcal{M}) \subset C^{0,\frac12}([0,1];\mathcal{M})$ and thus Hölder continuity of paths with finite path energy. Now, let $\hat y^K$ denote the piecewise geodesic interpolation of $(y(\frac{0}{K}),\ldots,y(\frac{K}{K}))$. For any K we have

$$\begin{aligned}\mathcal{E}[(y(t))_{t\in[0,1]}] &\geq K \textstyle\sum_{k=1}^K \mathrm{dist}^2(y(\frac{k-1}{K}),y(\frac{k}{K}))\\ &\geq K \textstyle\sum_{k=1}^K \mathcal{W}[y(\frac{k-1}{K}),y(\frac{k}{K})] - C\,K\,\sum_{k=1}^K \mathrm{dist}^3(y(\frac{k-1}{K}),y(\frac{k}{K})))\\ &\geq \mathrm{E}^K[(\hat y^K(t))_{t\in[0,1]}] - C\,K\,\textstyle\sum_{k=1}^K K^{-\frac32}\mathcal{E}[(\hat y^K(t))_{t\in[\frac{k-1}{K},\frac{k}{K}]}]^{\frac32}\\ &\geq \mathrm{E}^K[(\hat y^K(t))_{t\in[0,1]}]\big(1 - C\,K^{-\frac12}\sqrt{\mathcal{E}[(y(t))_{t\in[0,1]}]}\big)\,,\end{aligned}$$

where we have used (5.6). Letting $K\to\infty$ yields the desired limsup–inequality.
[*liminf-inequality*] We have to show that for any sequence of curves $(y^K)_{K\geq 1}$ with $y^K : [0,1]\to\mathcal{M}$ and $y^K\to y$ in $L^2((0,1);\mathbf{Y})$ the inequality $\liminf_{K\to\infty}\mathrm{E}^K[(y^K(t))_{t\in[0,1]}] \geq \mathcal{E}[(y(t))_{t\in[0,1]}]$ holds. Without any restriction we assume that $\mathrm{E}^K[(y^K(t))_{t\in[0,1]}] < \bar{\mathrm{E}} < \infty$ uniformly. Thus, $\mathcal{W}[y^K(\frac{k-1}{K}),y^K(\frac{k}{K})] \to 0$ uniformly as $K\to\infty$ so that due to the coercivity $\mathrm{dist}(y^K(\frac{k-1}{K}),y^K(\frac{k}{K}))\to 0$ uniformly as well. Next, we estimate

$$\begin{aligned}&\liminf_{K\to\infty}\mathrm{E}^K[(y^K(t))_{t\in[0,1]}] = \liminf_{K\to\infty} K\textstyle\sum_{k=1}^K\mathcal{W}[y^K(\frac{k-1}{K}),y^K(\frac{k}{K})]\\ &\quad\geq \liminf_{K\to\infty} K\textstyle\sum_{k=1}^K \mathrm{dist}^2(y^K(\frac{k-1}{K}),y^K(\frac{k}{K}))\,\big(1 - C\,\mathrm{dist}(y^K(\frac{k-1}{K}),y^K(\frac{k}{K}))\big)\\ &\quad= \liminf_{K\to\infty}\mathrm{E}[(y^K(t))_{t\in[0,1]}]\,,\end{aligned}$$

which also shows that y^K is uniformly bounded in $W^{1,2}((0,1);\mathcal{M})$. Due to the reflexivity of $\mathbf{V}$, a subsequence (for simplicity again denoted by $(y^K)_{K\geq1}$) weakly converges against y in $W^{1,2}((0,1);\mathcal{M})$. Due to the sequential weak lower semi-continuity of the energy $\mathcal{E}$ (*cf*. Theorem 5.3) we finally obtain the requested estimate

$$\liminf_{K\to\infty}\mathrm{E}^K[(y^K(t))_{t\in[0,1]}] \geq \liminf_{K\to\infty}\mathcal{E}[(y^K(t))_{t\in[0,1]}] \geq \mathcal{E}[(y(t))_{t\in[0,1]}]\,,$$

which concludes the proof. □

COROLLARY 5.10 (Convergence of discrete geodesics). *Minimizers of the discrete path energies* E^K *converge against minimizers of the continuous path energy* $\mathcal{E}$ *in* $C^0([0,1];\mathbf{Y})$.

Proof. This is a simple implication of the Γ-convergence and the following equi-mild coercivity of the discrete energies: Theorem 5.5 shows that the minima of the discrete energies are uniformly bounded. However, E^K is coercive with respect to $y_A + W^{1,2}((0,1);\mathbf{V})$ so that the minima for all K are achieved in some bounded ball of $y_A + W^{1,2}((0,1);\mathbf{V})$, which is compact in $L^2((0,1);\mathbf{Y})$. Together with the Γ-convergence of E^K, this implies that any $L^2((0,1);\mathbf{Y})$-limit point of a sequence of minimizers $\hat y^K$ for E^K is a minimizer for $\mathcal{E}$. Furthermore, from the above, all converging subsequences are bounded in $y_A + W^{1,2}((0,1);\mathbf{V})$ and thus bounded in $y_A + C^{0,\frac12}([0,1];\mathbf{V})$ and by Arzelà–Ascoli precompact in $C^0([0,1];\mathbf{Y})$ so that the convergence is not only in $L^2((0,1);\mathbf{Y})$, but even in $C^0([0,1];\mathbf{Y})$. □ The above convergence obviously does not only hold for the piecewise geodesic interpolation $\hat y^K$ of discrete geodesics $(y_0,\ldots,y_K)$, but also for the piecewise linear interpolation which we shall call y_τ (where $\tau = 1/K$ stands for the discrete time step).

For stronger convergence estimates and for the convergence of discrete logarithm, exponential map, and parallel transport, we require the following smoothness hypotheses:

(H1) The metric g is $C^2(\mathbf{Y}; \mathbf{V}' \otimes \mathbf{V}')$-smooth.
(H2) The energy $\mathcal{W}$ is $C^4((y_A + \mathbf{V}) \times (y_A + \mathbf{V}); \mathbb{R})$-smooth with bounded derivatives.

The following theorem now states that the convergence in $C^0([0,1]; \mathbf{Y})$ ensured by the above Γ-convergence result is actually much stronger with L^2-converging velocities.

THEOREM 5.11 (Path convergence in $W^{1,2}((0,1); \mathcal{M})$). *Under the hypotheses (H1) and (H2), if the interpolated geodesics $y_\tau(\cdot)$ converge for $K \to \infty$ to the continuous one $y(\cdot)$ in $L^2((0,1); \mathbf{Y})$, then this convergence is even in $W^{1,2}((0,1); \mathcal{M})$.*

Proof. Under the smoothness and coercivity assumptions on g, a continuous geodesic path $(y(t))_{t\in[0,1]}$ in $\mathcal{M}$ as a minimizer of the continuous energy $\int_0^1 g_{y(t)}(\dot{y}(t), \dot{y}(t))\,\mathrm{d}t$ for given fixed end points $y(0)$ and $y(1)$ is smooth and fulfills the Euler–Lagrange equation

$$\int_0^1 2g_{y(t)}(\dot{y}(t), \dot{\psi}(t)) + \big(D_y g_{y(t)}(\psi(t))\big)\,(\dot{y}(t), \dot{y}(t))\,\mathrm{d}t = 0 \tag{5.7}$$

for all $\psi \in W_0^{1,2}((0,1); \mathbf{V}) := \{\psi \in W^{1,2}((0,1); \mathbf{V}) \,|\, \psi(0) = \psi(1) = 0\}$. Furthermore, the Euler–Lagrange equation for a discrete geodesic path is given by

$$K \textstyle\sum_{k=1}^K \left(\mathcal{W}_{,1}[y_{k-1}, y_k](\psi_{k-1}) + \mathcal{W}_{,2}[y_{k-1}, y_k](\psi_k)\right) = 0 \tag{5.8}$$

for all $(\psi_k)_{k=0,\ldots,K} \subset \mathbf{V}$ with $\psi(0) = \psi(K) = 0$. Applying the Taylor expansion

$$\begin{aligned} \mathcal{W}[y_{k-1}, y_k] &= \mathcal{W}[y_{k-1}, y_{k-1}] + \mathcal{W}_{,2}[y_{k-1}, y_{k-1}](y_k - y_{k-1}) \\ &\quad + \int_0^1 (1-s)\mathcal{W}_{,22}[y_{k-1}, y_{k-1} + s(y_k - y_{k-1})](y_k - y_{k-1}, y_k - y_{k-1})\,\mathrm{d}s\,, \end{aligned} \tag{5.9}$$

whose first two terms on the right-hand side vanish, one can rewrite (5.8) as

$$\begin{aligned} 0 = \frac{1}{\tau}\sum_{k=1}^K \int_0^1 (1-s)\Big(& W_{,221}[y_{k-1}, y_{k-1}+s(y_k-y_{k-1})]((y_k-y_{k-1}), (y_k-y_{k-1}), \psi_{k-1}) \\ & +W_{,222}[y_{k-1}, y_{k-1}+s(y_k-y_{k-1})] \\ & \qquad ((y_k-y_{k-1}), (y_k-y_{k-1}), \psi_{k-1} + s(\psi_k-\psi_{k-1})) \\ & +2\,W_{,22}[y_{k-1}, y_{k-1}+s(y_k-y_{k-1})]((y_k-y_{k-1}), (\psi_k-\psi_{k-1}))\Big)\;\mathrm{d}s \\ = \sum_{k=1}^K \int_0^\tau (1-t/\tau)\Big(& W_{,221}[y_{k-1}, y_{k-1}+tv_k](v_k, v_k, \psi_{k-1}) \\ & +W_{,222}[y_{k-1}, y_{k-1}+tv_k](v_k, v_k, \psi_{k-1} + tw_k) \\ & +2\,W_{,22}[y_{k-1}, y_{k-1}+tv_k](v_k, w_k)\Big)\;\mathrm{d}t \end{aligned}$$

for $v_k = \frac{y_k - y_{k-1}}{\tau}$ and $w_k = \frac{\psi_k - \psi_{k-1}}{\tau}$. Now, taking into account the smoothness of the geodesic, the smoothness of $\mathcal{W}$, and $g_y(v,w) = \frac{1}{2}\mathcal{W}_{,22}[y,y](v,w)$, which implies

$$2(D_y g_y(\psi))(v,w) = \left(\mathcal{W}_{,221}[y,y](v,w,\psi) + \mathcal{W}_{,222}[y,y](v,w,\psi)\right)\,, \tag{5.10}$$

we finally achieve

$$\begin{aligned} 0 &= \frac{1}{2}\sum_{k=1}^{K}\tau\Big(W_{,221}[y_{k-1},y_{k-1}](v_k,v_k,\psi_{k-1}) + W_{,222}[y_{k-1},y_{k-1}](v_k,v_k,\psi_{k-1}) \\ &\qquad\qquad +2\,W_{,22}[y_{k-1},y_{k-1}](v_k,w_k)\Big) \ + \mathrm{Err} \\ &= \sum_{k=1}^{K}\tau\Big(2g_{y_{k-1}}(v_k,w_k) + \Big((D_y g_{y_{k-1}})(\psi_{k-1})\Big)(v_k,v_k)\Big) + \mathrm{Err} \end{aligned} \tag{5.11}$$

for $\mathrm{Err} = O(\tau^2 \sum_k \|v_k\|_{\mathbf{V}}^3 \|\psi_{k-1}\|_{\mathbf{V}}) + O(\tau^2 \sum_k \|v_k\|_{\mathbf{V}}^2 \|w_k\|_{\mathbf{V}})$. From Theorem 5.6 we obtain the following estimates for the error,

$$\mathrm{Err} = O(\tau^2)\left(\textstyle\sum_{k=1}^{K}\|\psi_{k-1}\|_{\mathbf{V}} + \sum_{k=1}^{K}\|w_k\|_{\mathbf{V}}\right), \tag{5.12}$$

$$\mathrm{Err} = O(\tau)\left(\sqrt{\tau\textstyle\sum_{k=1}^{K}\|\psi_{k-1}\|_{\mathbf{V}}^2} + \sqrt{\tau\textstyle\sum_{k=1}^{K}\|w_k\|_{\mathbf{V}}^2}\right). \tag{5.13}$$

Based on these preliminaries we now combine the Euler–Lagrange equation (5.7) for the continuous geodesic path and (5.11) (derived from the Euler–Lagrange equation (5.8) for the discrete geodesic path) to obtain an equation for the discretization error. To this end, we consider a piecewise polygonal function $\psi_\tau : [0,1] \to \mathbf{V}$ with $\psi_\tau(k\tau) = \psi_k \in \mathbf{V}$ for $k = 0,\dots,K$ and $\psi_0 = \psi_K = 0$. Using this notation one easily verifies the identity

$$\begin{aligned} &\textstyle\int_0^1 2g_{y(t)}(\dot y_\tau(t) - \dot y(t), \dot\phi(t))\,\mathrm{d}t \\ &= \textstyle\int_0^1 2g_{y(t)}(\dot y_\tau(t) - \dot y(t), \dot\phi(t) - \dot\psi_\tau(t))\,\mathrm{d}t \\ &\quad -\textstyle\int_0^1 2g_{y(t)}(\dot y(t), \dot\psi_\tau(t)) + D_y g_{y(t)}(\psi_\tau(t))(\dot y(t), \dot y(t))\,\mathrm{d}t \\ &\quad +\textstyle\sum_{k=1}^K \tau\Big(2g_{y_{k-1}}(v_k,w_k) + D_y g_{y_{k-1}}(\psi_{k-1})(v_k,v_k)\Big) \\ &\quad +\textstyle\sum_{k=1}^K 2\left(\int_{(k-1)\tau}^{k\tau} g_{y_\tau(t)}(\dot y_\tau(t), \dot\psi_\tau(t))\,\mathrm{d}t - \tau g_{y_{k-1}}(v_k,w_k)\right) \\ &\quad +2\textstyle\int_0^1 g_{y(t)}(\dot y_\tau(t), \dot\psi_\tau(t)) - g_{y_\tau(t)}(\dot y_\tau(t), \dot\psi_\tau(t))\,\mathrm{d}t \\ &\quad +\textstyle\sum_{k=1}^K \left(\int_{(k-1)\tau}^{k\tau} D_y g_{y_\tau(t)}(\psi_{k-1})(\dot y_\tau(t), \dot y_\tau(t))\,\mathrm{d}t - \tau D_y g_{y_{k-1}}(\psi_{k-1})(v_k,v_k)\right) \\ &\quad +\textstyle\sum_{k=1}^K \left(\int_{(k-1)\tau}^{k\tau} D_y g_{y_\tau(t)}(\psi_\tau(t))(v_k,v_k) - D_y g_{y_\tau(t)}(\psi_{k-1})(\dot y_\tau(t), \dot y_\tau(t))\,\mathrm{d}t\right) \\ &\quad +\textstyle\int_0^1 \left(D_y g_{y(t)}(\psi_\tau(t)) - D_y g_{y_\tau(t)}(\psi_\tau(t))\right)(\dot y_\tau(t), \dot y_\tau(t))\,\mathrm{d}t \\ &\quad +\textstyle\int_0^1 D_y g_{y(t)}(\psi_\tau(t))(\dot y_\tau(t), \dot y(t) - \dot y_\tau(t)) + D_y g_{y(t)}(\psi_\tau(t))(\dot y(t) - \dot y_\tau(t), \dot y(t))\,\mathrm{d}t \\ &=: I - II + III + IV + V + VI + VII + VIII + IX \end{aligned} \tag{5.14}$$

with $v(t) = \dot y(t)$, $w(t) = \dot\psi_\tau(t)$. Now, we choose $\phi = y_\tau - y$ and $\psi_\tau = y_\tau - \mathcal{I}_\tau y$, where $\mathcal{I}_\tau$ is the piecewise affine Lagrangian interpolation in time with $\mathcal{I}_\tau\psi(k\tau) = \psi(k\tau)$. Let us mention here, that we will reuse (5.14) with different test functions later in the context of a pointwise error estimate in the proof of Theorem 6.1. Due to the uniform coercivity of the metric g the left-hand side can be estimated from below by $2c^*\|\dot y - \dot y_\tau\|_{L^2((0,1);\mathbf{V})}^2$. The different terms on the right-hand side of (5.14) are estimated as follows:

- (I) By the uniform boundedness of the metric, $|\, I\,|$ can be estimated by $2C^*\|\dot y - \dot y_\tau\|_{L^2((0,1);\mathbf{V})}\|\dot y - (\mathcal{I}_\tau y)^\cdot\|_{L^2((0,1);\mathbf{V})}$.
- (II) Due to (5.7) the term II vanishes.

(III) Taking into account (5.11) and (5.13), $|\,III\,|$ is bounded by $C\tau\|\psi_\tau\|_{W^{1,2}((0,1);\mathbf{V})}$.

(IV) Due to the smoothness of the metric g and Theorem 5.6, the quadrature error in $|\,IV\,|$ is bounded by $O(\tau^2\sum_k\|v_k\|_{\mathbf{V}}^2\|\dot\psi_\tau\|_{\mathbf{V}}) = O(\tau\|\dot\psi_\tau\|_{L^2((0,1);\mathbf{V})})$ (compare (5.13)).

(V) By an analogous argument and the boundedness of y_τ in $L^2((0,1);\mathbf{Y})$ the term $|\,V\,|$ can be estimated by $C\|y-y_\tau\|_{L^2((0,1);\mathbf{Y})}\|\dot y_\tau\|_{L^\infty((0,1);\mathbf{V})}\|\dot\psi_\tau\|_{L^2((0,1);\mathbf{V})}$, where we know from Theorem 5.6 that $\|\dot y_\tau\|_{L^\infty((0,1);\mathbf{V})} = O(1)$.

(VI) Now using the smoothness of $D_y g$ we can bound $|\,VI\,|$ by $C\tau^2\sum_{k=1}^K\|v_k\|_{\mathbf{V}}^2\|v_k\|_{\mathbf{Y}}\|\psi_{k-1}\|_{\mathbf{Y}} = O(\tau\|\psi_\tau\|_{L^2((0,1);\mathbf{Y})})$.

(VII) Term $|\,VII\,|$ is bounded by $C\tau\|\dot\psi_\tau\|_{L^2((0,1);\mathbf{Y})}$ due to the uniform bound for $\|\dot y_\tau\|_{\mathbf{V}}$.

(VIII) Now using the smoothness of $D_y g$ and the boundedness of $\|\dot y_\tau\|_{\mathbf{V}}$ we can bound $|\,VIII\,|$ by $C\|y-y_\tau\|_{L^2((0,1);\mathbf{Y})}\|\psi_\tau\|_{L^2((0,1);\mathbf{Y})}$.

(IX) Finally, we obtain the estimate $|\,IX\,| \leq C\|\psi_\tau\|_{L^2((0,1);\mathbf{Y})}\|\dot y-\dot y_\tau\|_{L^2((0,1);\mathbf{V})}$.

Altogether, for some constant C we obtain the following estimate for the error,

$$\begin{aligned}2c^*\|\dot y-\dot y_\tau\|^2_{L^2((0,1);\mathbf{V})} \leq{}& 2C^*\|\dot y-\dot y_\tau\|_{L^2((0,1);\mathbf{V})}\|\dot y-(\mathcal{I}_\tau y)^\cdot\|_{L^2((0,1);\mathbf{V})}\\ &+C\tau\|\psi_\tau\|_{W^{1,2}((0,1);\mathbf{V})}\\ &+C\|y-y_\tau\|_{L^2((0,1);\mathbf{Y})}\|\dot\psi_\tau\|_{L^2((0,1);\mathbf{V})}\\ &+C\tau\|\psi_\tau\|_{W^{1,2}((0,1);\mathbf{Y})}\\ &+C\|y-y_\tau\|_{L^2((0,1);\mathbf{Y})}\|\psi_\tau\|_{L^2((0,1);\mathbf{Y})}\\ &+C\|\psi_\tau\|_{L^2((0,1);\mathbf{Y})}\|\dot y-\dot y_\tau\|_{L^2((0,1);\mathbf{V})}\,.\end{aligned}$$

Finally, using Poincaré's inequality $\|y-y_\tau\|_{L^2((0,1);\mathbf{V})} \leq \|\dot y-\dot y_\tau\|_{L^2((0,1);\mathbf{V})}$ as well as Young's inequality and the compact embedding $\mathbf{V}\hookrightarrow\mathbf{Y}$, we derive in a straightforward way

$$\|\dot y-\dot y_\tau\|^2_{L^2((0,1);\mathbf{V})} \leq C\left(\tau^2+\|y-\mathcal{I}_\tau y\|^2_{W^{1,2}((0,1);\mathbf{V})}+\|y-y_\tau\|^2_{L^2((0,1);\mathbf{Y})}\right)$$

for some $C>0$. Taking into account the classical interpolation estimate

$$\|y-\mathcal{I}_\tau y\|_{L^2((0,1);\mathbf{V})}+\tau\|\dot y-(\mathcal{I}_\tau y)^\cdot\|_{L^2((0,1);\mathbf{V})} \leq C\tau^2$$

as well as $\|y-y_\tau\|_{L^2((0,1);\mathbf{Y})}\to 0$, this concludes the proof. □

REMARK 5.12. *Note that in the continuous setting, geodesics can also be obtained by minimizing the path length* (2.1) *for fixed end points* $y(0), y(1)$*. However, discrete minimizers of the discrete path energy* (2.1) *are in general unrelated to continuous geodesics. As an example, think of a smooth curved manifold embedded in* $\mathbb{R}^m$ *and let* $\mathcal{W}[y,\tilde y]$ *be the squared Euclidean distance between* y *and* $\tilde y$ *in the embedding space. Then given* y_0, y_K *with* $y_0+t(y_1-y_0)\notin\mathcal{M}$ *for all* $t\in(0,1)$*, minimizers of* $\mathrm{L}[\cdot]$ *for* y_0, y_K *fixed have the form* $(y_0,y_0,\ldots,y_0,y_K,\ldots,y_K)$*. Obviously, the claim of Theorem* 5.6 *no longer holds, and the argument for Theorem* 5.9 *breaks down since minimizers of the discrete length* $\mathrm{L}[\cdot]$ *are only bounded in* $BV((0,1);\mathcal{M})$ *instead of* $W^{1,2}((0,1);\mathcal{M})$*.*

6. Convergence of discrete logarithm, exponential, and parallel transport. In this section we discuss the limit behavior of the discrete operators. At first we investigate the convergence of the discrete logarithm, which can be formulated as an L^∞ derivative error estimate for discrete variational solutions of an elliptic problem in $W^{1,2}((0,1);\mathcal{M})$. This will become apparent in the proof of the following theorem.

THEOREM 6.1 (Convergence of discrete logarithm). *Let* $y,\tilde y\in\mathcal{M}$*. Under the hypotheses (H1) and (H2), if the continuous and discrete geodesics between* $y,\tilde y$ *are unique,* $K(\frac{1}{K}\mathrm{LOG})_y\tilde y\to\log_y\tilde y$ *weakly in* $\mathbf{V}$ *(and thus strongly in* $\mathbf{Y}$*) as* $K\to\infty$*.*

Proof. As usual, abbreviate $\tau = 1/K$, and let $y(t)_{t\in[0,1]}$ and $y_\tau(t)_{t\in[0,1]}$ be the continuous and the interpolated discrete geodesic between y and $\tilde y$. Let us denote by $\Phi \in W_0^{1,2}((0,1);\mathrm{Hom}(\mathbf{V}',\mathbf{V}))$ the weak solution of

$$\int_0^1 2g_{y(t)}(\dot z(t), \dot\Phi(t)\mathfrak{v})\,\mathrm{d}t = \left(\mathfrak{v}, \tfrac{z(\tau)}{\tau}\right)_{\mathbf{V}',\mathbf{V}} \tag{6.1}$$

for all $z \in W_0^{1,2}((0,1);\mathbf{V})$, $\mathfrak{v} \in \mathbf{V}'$. Then we obtain

$$\begin{aligned}\left(\mathfrak{v}, K(\tfrac{1}{K}\mathrm{LOG})_y\tilde y - \log_y \tilde y\right)_{\mathbf{V}',\mathbf{V}} &= \left(\mathfrak{v}, \tfrac{y_\tau(\tau)-y(0)}{\tau} - \dot y(0)\right)_{\mathbf{V}',\mathbf{V}}\\ &= \left(\mathfrak{v}, \tfrac{y_\tau(\tau)-y(\tau)}{\tau} + \left(\tfrac{y(\tau)-y(0)}{\tau} - \dot y(0)\right)\right)_{\mathbf{V}',\mathbf{V}}\\ &= \int_0^1 2g_{y(t)}(\dot y_\tau(t) - \dot y(t), \dot\Phi(t)\mathfrak{v})\,\mathrm{d}t + O(\tau\|\mathfrak{v}\|_{\mathbf{V}'})\,. \end{aligned} \tag{6.2}$$

The solution of (6.1) for a given continuous geodesic path $(y(t))_{t\in[0,1]}$ can be computed explicitly. Indeed, we deduce from (6.1) the following two conditions,

$$2\frac{\mathrm{d}}{\mathrm{d}t}\left(\mathscr{G}_{y(t)}(\dot\Phi(t)\mathfrak{v})\right) = 0 \quad \text{on } (0,\tau)\cup(\tau,1)\,, \tag{6.3}$$

$$2\,\mathscr{G}_{y(\tau)}\left(\dot\Phi(\tau+0)\mathfrak{v} - \dot\Phi(\tau-0)\mathfrak{v}\right) = -\tfrac{1}{\tau}\mathfrak{v} \tag{6.4}$$

for any $\mathfrak{v} \in \mathbf{V}'$, where we use the notation $\mathscr{G}_y : \mathbf{V} \to \mathbf{V}'$ for the inverse Riesz isomorphism of the Hilbert space $(\mathbf{V}, g_y)$. Integration of (6.3) leads to $\dot\Phi(t) = \mathscr{G}_{y(t)}^{-1}\mathscr{G}_{y(0)}\dot\Phi(0)$ on $(0,\tau)$ and $\dot\Phi(t) = \mathscr{G}_{y(t)}^{-1}\mathscr{G}_{y(1)}\dot\Phi(1)$ on $(\tau,1)$. From this and (6.4) one obtains $\mathscr{G}_{y(1)}\dot\Phi(1) - \mathscr{G}_{y(0)}\dot\Phi(0) = -\frac{1}{2\tau}$. Furthermore, due to the boundary conditions $\Phi(0) = \Phi(1) = 0$ we achieve

$$0 = \int_0^\tau \dot\Phi(t)\,\mathrm{d}t + \int_\tau^1 \dot\Phi(t)\,\mathrm{d}t = \left(\int_0^\tau \mathscr{G}_{y(t)}^{-1}\,\mathrm{d}t\right)\mathscr{G}_{y(0)}\dot\Phi(0) + \left(\int_\tau^1 \mathscr{G}_{y(t)}^{-1}\,\mathrm{d}t\right)\mathscr{G}_{y(1)}\dot\Phi(1) \tag{6.5}$$

Now, with the notation $G^- = \int_0^\tau \mathscr{G}_{y(t)}^{-1}\,\mathrm{d}t$, $G^+ = \int_\tau^1 \mathscr{G}_{y(t)}^{-1}\,\mathrm{d}t$, $A = \mathscr{G}_{y(0)}\dot\Phi(0)$, and $B = \mathscr{G}_{y(1)}\dot\Phi(1)$, (6.4) and (6.5) form the linear system of equations

$$\begin{aligned} A - \quad B &= \tfrac{1}{2\tau}\\ G^-A + G^+B &= 0 \end{aligned}$$

which for $\mathbb{1}$ the identity has the unique solution

$$A = \frac{1}{2\tau}(G^-)^{-1}G^+(\mathbb{1} + (G^-)^{-1}G^+)^{-1}\,, \quad B = -\frac{1}{2\tau}(\mathbb{1} + (G^-)^{-1}G^+)^{-1}\,.$$

Finally, we can evaluate $\Phi(t)$ via integration of

$$\dot\Phi(t) = \frac{1}{2\tau}\mathscr{G}_{y(t)}^{-1}\left(\int_0^\tau \mathscr{G}_{y(s)}^{-1}\,\mathrm{d}s\right)^{-1}\int_\tau^1 \mathscr{G}_{y(s)}^{-1}\,\mathrm{d}s\left(\mathbb{1} + \left(\int_0^\tau \mathscr{G}_{y(s)}^{-1}\,\mathrm{d}s\right)^{-1}\int_\tau^1 \mathscr{G}_{y(s)}^{-1}\,\mathrm{d}s\right)^{-1} \quad \text{on } (0,\tau)\,,$$

$$\dot\Phi(t) = -\frac{1}{2\tau}\mathscr{G}_{y(t)}^{-1}\left(\mathbb{1} + \left(\int_0^\tau \mathscr{G}_{y(s)}^{-1}\,\mathrm{d}s\right)^{-1}\int_\tau^1 \mathscr{G}_{y(s)}^{-1}\,\mathrm{d}s\right)^{-1} \quad \text{on } (\tau,1)\,.$$

Let us remark that in the trivial case of the constant metric $g_y(v,v) = (v,v)_{\mathbf{V}}$ on a Hilbert space $\mathbf{V}$ (in which case geodesics are straight lines), Φ is a piecewise affine function with $\Phi(0) = \Phi(1) = 0$ and $\Phi(\tau) = \frac{1-\tau}{2}\mathscr{R}$ for the Riesz isomorphism $\mathscr{R} : \mathbf{V}' \to \mathbf{V}$.

Next, we define Ψ_τ as the Lagrangian interpolation $\mathcal{I}_\tau\Phi$ of Φ with $\Psi_\tau(k\tau) = \Phi(k\tau)$ for $k = 0, \ldots, K$. To show the claim, it remains to show the convergence of the right-hand side in (6.2). At first we proceed as in the proof of Theorem 5.11. In (5.14) we choose $\phi = \Phi\mathfrak{v}$ and $\psi_\tau = \Psi_\tau\mathfrak{v}$, which turns the left-hand side into the desired error representation. Hence, it remains to verify that the terms I to IX on the right-hand side of (5.14) vanish for $\tau \to 0$. Indeed, using the same notation as in Theorem 5.11 we obtain the following estimates:

(I) Due to the uniform boundedness of the metric, $|\,I\,|$ can be bounded by $2C^*\|\dot y - \dot y_\tau\|_{L^2((0,1);\mathbf{V})}\|(\Phi - \mathcal{I}_\tau\Phi)^{\boldsymbol{\cdot}}\mathfrak{v}\|_{L^2((0,1);\mathbf{V})}$, and for $t \in ((k-1)\tau, k\tau)$ one obtains

$$\begin{aligned}\|(\Phi - \mathcal{I}_\tau\Phi)^{\boldsymbol{\cdot}}(t)\mathfrak{v}\|_{\mathbf{V}} &= \|\tfrac{1}{\tau}\textstyle\int_{(k-1)\tau}^{k\tau}(\dot\Phi(t) - \dot\Phi(s))\mathfrak{v}\;\mathrm{d}s\|_{\mathbf{V}}\\ &\le \tfrac{1}{\tau}\textstyle\int_{(k-1)\tau}^{k\tau}\|\mathscr{G}_{y(t)}^{-1} - \mathscr{G}_{y(s)}^{-1}\|\;\mathrm{d}s\,\|\mathscr{G}_{y(\hat t)}\dot\Phi(\hat t)\mathfrak{v}\|_{\mathbf{V}} = O(\tau)\|\dot\Phi(\hat t)\mathfrak{v}\|_{\mathbf{V}}\end{aligned}$$

with $\hat t = 0$ for $k = 1$ and $\hat t = 1$ else. Using $\dot\Phi(0) = O(\tau^{-1})$ and $\dot\Phi(1) = O(1)$, we obtain $\|(\Phi - \mathcal{I}_\tau\Phi)^{\boldsymbol{\cdot}}\mathfrak{v}\|_{L^2((0,1);\mathbf{V})} = O(\tau\|\mathfrak{v}\|_{\mathbf{V}'})$.

(II) Due to (5.7) the term II vanishes as before.

(III) By (5.11) and this time (5.12) the term $|\,III\,|$ can be estimated by

$$C\tau\|\psi_\tau\|_{L^\infty((0,1);\mathbf{V})} + C\tau^2\|w_1\|_{\mathbf{V}} + C\tau\sup_{k>1}\|w_k\|_{\mathbf{V}}\,,$$

where $\|w_1\|_{\mathbf{V}} \le C\tau^{-1}\|\mathfrak{v}\|_{\mathbf{V}'}$ and $\sup_{k>1}\|w_k\|_{\mathbf{V}} = O(\|\mathfrak{v}\|_{\mathbf{V}'})$.
Using $\|\psi_\tau\|_{L^\infty((0,1);\mathbf{V})} = O(1)$, $|\,III\,|$ is bounded by $C\tau\|\mathfrak{v}\|_{\mathbf{V}'}$.

(IV) Due to the smoothness of the metric g, the quadrature error in IV is bounded by

$$O(\tau^2\sum_k\|v_k\|_{\mathbf{Y}}\|v_k\|_{\mathbf{V}}\|\dot\psi_\tau\|_{\mathbf{V}}) = O(\tau^2\|w_1\|_{\mathbf{V}} + \tau\sup_{k>1}\|w_k\|_{\mathbf{V}}) = O(\tau\|\mathfrak{v}\|_{\mathbf{V}'})\,.$$

(V) Using Theorem 5.6, one obtains

$$\begin{aligned}|\,V\,| &\le C\textstyle\int_0^1\|y - y_\tau\|_{\mathbf{Y}}\|\dot\psi_\tau\|_{\mathbf{V}}\|\dot y_\tau\|_{\mathbf{V}}\;\mathrm{d}t\\ &\le C\|y - y_\tau\|_{L^\infty((0,1);\mathbf{Y})}\|\dot\psi_\tau\|_{L^1((0,1);\mathbf{V})}\|\dot y_\tau\|_{L^\infty((0,1);\mathbf{V})}\\ &= O(\|y - y_\tau\|_{L^\infty((0,1);\mathbf{Y})}\|\mathfrak{v}\|_{\mathbf{V}'})\,.\end{aligned}$$

(VI) By the smoothness of g we get $|\,VI\,| \le C\tau^2\sum_k\|\psi_{k-1}\|_{\mathbf{Y}}\|v_k\|_{\mathbf{V}}^2\|v_k\|_{\mathbf{Y}} = O(\tau\|\mathfrak{v}\|_{\mathbf{V}'})\,.$

(VII) Again due to the smoothness of g we achieve

$$|\,VII\,| \le C\tau^2\|w_1\|_{\mathbf{Y}}\|v_1\|_{\mathbf{V}}^2 + C\sum_{k>1}\tau^2\|w_k\|_{\mathbf{Y}}\|v_k\|_{\mathbf{V}}^2 \le C\tau\|\mathfrak{v}\|_{\mathbf{V}'}\,.$$

(VIII) For the term $VIII$ one obtains the estimate

$$\begin{aligned}|\,VIII\,| &\le C\|y - y_\tau\|_{L^2((0,1);\mathbf{Y})}\|\psi_\tau\|_{L^2((0,1);\mathbf{Y})}\|\dot y_\tau\|_{L^\infty((0,1);\mathbf{V})}^2\\ &= O(\|y - y_\tau\|_{L^2((0,1);\mathbf{Y})}\|\mathfrak{v}\|_{\mathbf{V}'})\,.\end{aligned}$$

(IX) Finally, for the last term we get

$$\begin{aligned}|\,IX\,| &\le C\|\psi_\tau\|_{L^2((0,1);\mathbf{Y})}(\|\dot y\|_{L^\infty((0,1);\mathbf{V})} + \|\dot y_\tau\|_{L^\infty((0,1);\mathbf{V})})\|\dot y - \dot y_\tau\|_{L^2((0,1);\mathbf{V})}\\ &= O(\|\dot y - \dot y_\tau\|_{L^2((0,1);\mathbf{V})}\|\mathfrak{v}\|_{\mathbf{V}'})\,.\end{aligned}$$

Collecting all these estimates we obtain

$$\Big(\mathfrak{v}, K(\tfrac{1}{K}\mathrm{LOG})_y\tilde y - \log_y \tilde y\Big)_{\mathbf{V}',\mathbf{V}} \le C\big(\tau + \|\dot y - \dot y_\tau\|_{L^2((0,1);\mathbf{V})} + \|y - y_\tau\|_{L^\infty((0,1);\mathbf{Y})}\big)\|\mathfrak{v}\|_{\mathbf{V}'}\,.$$

Using Theorems 5.10 and 5.11 the right hand side of this estimate converges to 0 as $K \to \infty$ for all $\mathfrak{v} \in \mathbf{V}'$, which proves the claim. $\square$

Next we consider the existence and convergence of the discrete exponential. This requires several preparations.

LEMMA 6.2 (Local uniqueness of $(\frac{1}{2}\mathrm{LOG})$). *Under the hypotheses (H1) and (H2), there exists an $\epsilon > 0$ such that for any $y_0, y_2 \in \mathcal{M}$ with $y_2 \in B_\epsilon(y_0) = \{y \mid \|y - y_0\|_\mathbf{V} \le \epsilon\}$, $\mathrm{E}[(y_0,\cdot,y_2)] : \mathcal{M} \to \mathbb{R}$ is strictly convex with bounded coercive Hessian on $B_\epsilon(y_0)$, and $(\frac{1}{2}\mathrm{LOG})_{y_0}(y_2)$ is unique.*

Proof. For given ϵ, consider $y_0, y_2 \in \mathcal{M}$ with $\|y_2 - y_0\|_\mathbf{V} < \epsilon$. The Hessian of $\mathrm{E}[(y_0,\cdot,y_2)]$ at some y_1 in the ball $B_\epsilon(y_0)$ is given by $2(\mathcal{W}_{,22}[y_0,y_1] + \mathcal{W}_{,11}[y_1,y_2]) = 8g_{y_0} + O(\epsilon)$ with uniform constants, where we have used the smoothness of $\mathcal{W}$ and Lemma 5.7. For ϵ small enough, this is bounded and coercive (independent of y_0), and thus $\mathrm{E}[(y_0,\cdot,y_2)]$ is strictly convex (cf. Theorem 5.8). This also implies uniqueness of $(\frac{1}{2}\mathrm{LOG})_{y_0}(y_2)$ (via Theorem 5.5 it is easy to see that the optimal y_1 cannot lie outside $B_\epsilon(y_0)$). $\square$

REMARK 6.3. *The above result can be generalized to K-geodesics inside an ϵ-ball. It is a refinement of Theorem 5.8 in the sense that it provides a uniform uniqueness radius ϵ. Also, it implies the differentiability of $(\frac{1}{2}\mathrm{LOG})_{y_0}(y_2)$ with respect to $y_2 \in B_\epsilon(y_0)$: The Hessian $D^2\mathrm{E}[(y_0,\cdot,y_2)] = 2(\mathcal{W}_{,22}[y_0,\cdot] + \mathcal{W}_{,11}[\cdot,y_2])$ is coercive so that by the implicit function theorem,*

$$\begin{aligned} D(\tfrac{1}{2}\mathrm{LOG})_{y_0}(y_2) &= -\big(\mathcal{W}_{,22}[y_0,y_1] + \mathcal{W}_{,11}[y_1,y_2]\big)^{-1}\mathcal{W}_{,12}[y_1,y_2] \\ &= [4g_{y_0} + O(\epsilon)]^{-1}(2g_{y_0} + O(\epsilon)) = \tfrac{1}{2}\mathbb{1} + O(\epsilon) \end{aligned}$$

for $y_1 := y_0 + (\frac{1}{2}\mathrm{LOG})_{y_0}(y_2)$.

LEMMA 6.4. *For $y_2 \in B_\epsilon(y_0)$ the estimate $y_0 + (\frac{1}{2}\mathrm{LOG})_{y_0}(y_2) = \frac{y_0+y_2}{2} + O(\epsilon^{3/2})$ holds.*

Proof. Indeed, using $\mathcal{W}[y,\tilde y] = \mathrm{dist}^2(y,\tilde y) + O(\mathrm{dist}^3(y,\tilde y)) = g_{y_0}(y-\tilde y, y-\tilde y) + O(\epsilon^3)$ for $y, \tilde y \in B_\epsilon(y_0)$ and $2[\mathcal{W}[y_0, y_1(y_2)] + \mathcal{W}[y_1(y_2), y_2)]] = g_{y_0}(y_2 - y_0, y_2 - y_0) + O(\epsilon^3)$ for $y_1(y_2) = y_0 + (\frac{1}{2}\mathrm{LOG})_{y_0}(y_2)$ (by Theorem 5.5), we get

$$\begin{aligned} &2g_{y_0}\big(\tfrac{y_0+y_2}{2} - y_1(y_2), \tfrac{y_0+y_2}{2} - y_1(y_2)\big) \\ &\quad= g_{y_0}(y_0 - y_1(y_2), y_0 - y_1(y_2)) + g_{y_0}(y_2 - y_1(y_2), y_2 - y_1(y_2)) - \tfrac{1}{2}g_{y_0}(y_2 - y_0, y_2 - y_0) \\ &\quad= \mathcal{W}[y_0, y_1(y_2)] + \mathcal{W}[y_1(y_2), y_2] - (\mathcal{W}[y_0, y_1(y_2)] + \mathcal{W}[y_1(y_2), y_2]) + O(\epsilon^3) = O(\epsilon^3)\,. \square \end{aligned}$$

LEMMA 6.5 (Local existence of (EXP^2)). *Under the hypotheses (H1) and (H2), there exists an $\epsilon > 0$ such that $(\mathrm{EXP}^2)_{y_0}(v)$ exists for any $y_0 \in \mathcal{M}$, $v \in \mathbf{V}$, with $\|v\|_\mathbf{V} < \epsilon$.*

Proof. Choose ϵ to be one third the value from Lemma 6.2. For $y_0 \in \mathcal{M}$, $v \in \mathbf{V}$ with $\|v\|_\mathbf{V} < \epsilon$, define the operator $T_v : B_{3\epsilon}(y_0) \to \mathcal{M}$ by $T_v y_2 = y_2 + v - (\frac{1}{2}\mathrm{LOG})_{y_0}(y_2)$. Without loss of generality we may assume ϵ to be small enough such that by Lemma 6.4, $T_v : B_{3\epsilon}(y_0) \to B_{3\epsilon}(y_0)$. Its derivative is given by $\mathbb{1} - D(\frac{1}{2}\mathrm{LOG})_{y_0}(y_2) = \frac{1}{2}\mathbb{1} + O(\epsilon)$, hence T_v is a contraction, and by the Banach fixed point theorem has a unique fixed point $(\mathrm{EXP}^2)_{y_0}(v)$. $\square$

REMARK 6.6. *By the inverse function theorem, $(\mathrm{EXP}^2)_{y_0}$ is differentiable at v with $\|v\|_\mathbf{V} < \epsilon$, and $D(\mathrm{EXP}^2)_{y_0}(v) = [D(\frac{1}{2}\mathrm{LOG})_{y_0}((\mathrm{EXP}^2)_{y_0}(v))]^{-1} = 2\mathbb{1} + O(\epsilon)$.*

LEMMA 6.7. *For $\|v\|_\mathbf{V} \le \epsilon$ the estimate $(\mathrm{EXP}^2)_{y_0}(v) = (y_0 + 2v) + O(\epsilon^{3/2})$ holds.*

Proof. Let us abbreviate $y_1 = y_0 + v$ and $y_2(y_1) = (\mathrm{EXP}^2)_{y_0}(v)$. Then as in Lemma 6.4,
$\frac{1}{2} g_{y_0}(y_0 + 2v - y_2(y_1), y_0 + 2v - y_2(y_1))$
$= g_{y_0}(y_0 - y_1, y_0 - y_1) + g_{y_0}(y_2(y_1) - y_1, y_2(y_1) - y_1) - \frac{1}{2} g_{y_0}(y_2(y_1) - y_0, y_2(y_1) - y_0)$
$= \mathcal{W}[y_0, y_1] + \mathcal{W}[y_1, y_2(y_1)] - (\mathcal{W}[y_0, y_1] + \mathcal{W}[y_1, y_2(y_1)]) + O(\epsilon^3) = O(\epsilon^3)$. □

REMARK 6.8. *Using the above boundedness, our previous estimates can be further improved to* $\|(\mathrm{EXP}^2)_{y_0}(v) - (y_0 + 2v)\|_{\mathbf{V}} = O(\|v\|_{\mathbf{V}}^2)$ *and* $\|(\frac{1}{2}\mathrm{LOG})_{y_0}(y_2) - (\frac{y_0 + y_2}{2} - y_0)\|_{\mathbf{V}} = O(\|y_2 - y_0\|_{\mathbf{V}}^2)$ *for* $\|v\|_{\mathbf{V}}$ *and* $\|y_2 - y_0\|_{\mathbf{V}}$ *smaller than* ϵ*. Indeed, if* (y_0, y_1, y_2) *is a discrete geodesic with* $v_1 = v = y_1 - y_0$ *and* $v_2 = y_2 - y_1$*, and if* ψ_1 *is a variation of* y_1*, then the optimality condition implies*

$$\begin{aligned}
0 &= \mathcal{W}_{,2}[y_0, y_1](\psi_1) + \mathcal{W}_{,1}[y_1, y_2](\psi_1) \\
&= \mathcal{W}_{,2}[y_1, y_1](\psi_1) - \mathcal{W}_{,21}[y_1, y_1](\psi_1, y_1 - y_0) + O(\|y_1 - y_0\|_{\mathbf{V}}^2 \|\psi_1\|_{\mathbf{V}}) \\
&\quad + \mathcal{W}_{,1}[y_1, y_1](\psi_1) + \mathcal{W}_{,12}[y_1, y_1](\psi_1, y_2 - y_1) + O(\|y_2 - y_1\|_{\mathbf{V}}^2 \|\psi_1\|_{\mathbf{V}}) \\
&= 2 g_{y_1}(\psi_1, v_1 - v_2) + O((\|v_1\|_{\mathbf{V}}^2 + \|v_2\|_{\mathbf{V}}^2)\|\psi_1\|_{\mathbf{V}}) \,,
\end{aligned}$$

where we used Lemma 5.7. Using the coercivity of g_{y_1}*, the desired estimates can now easily be derived.*

THEOREM 6.9 (Existence and convergence of (EXP^K)). *Let* $y : [0, 1] \to \mathcal{M}$ *be a smooth geodesic. Under the hypotheses (H1) and (H2),* $(\mathrm{EXP}^K)_{y(0)}(\frac{\dot{y}(0)}{K})$ *exists for* K *large enough, and for* $\tau = \frac{1}{K}$ *one obtains* $\left\|y(1) - (\mathrm{EXP}^K)_{y(0)}(\frac{\dot{y}(0)}{K})\right\|_{\mathbf{V}} = O(\tau)$.

Proof. To examine the convergence of the discrete exponential, we need to linearize the optimality condition (5.8). Let $(y_0, \ldots, y_K)$ be some discrete path in $\mathcal{M}$ and let $v_k = (y_k - y_{k-1})/\tau$ for $\tau = 1/K$. As in the proof of Theorem 5.11, for $\psi_k \in \mathbf{V}$ we find

$$\begin{aligned}
&\mathcal{W}_{,2}[y_{k-1}, y_k](\psi_k) + \mathcal{W}_{,1}[y_k, y_{k+1}](\psi_k) \\
&= \tau \int_0^\tau (1 - \tfrac{t}{\tau}) \Big[\mathcal{W}_{,222}[y_{k-1}, y_{k-1} + t v_k](v_k, v_k, \tfrac{t}{\tau}\psi_k) + 2\mathcal{W}_{,22}[y_{k-1}, y_{k-1} + t v_k](v_k, \tfrac{1}{\tau}\psi_k) \\
&\quad + \mathcal{W}_{,221}[y_k, y_k + t v_{k+1}](v_{k+1}, v_{k+1}, \psi_k) + \mathcal{W}_{,222}[y_k, y_k + t v_{k+1}](v_{k+1}, v_{k+1}, \psi_k) \\
&\quad - \mathcal{W}_{,222}[y_k, y_k + t v_{k+1}](v_{k+1}, v_{k+1}, \tfrac{t}{\tau}\psi_k) - 2\mathcal{W}_{,22}[y_k, y_k + t v_{k+1}](v_{k+1}, \tfrac{1}{\tau}\psi_k) \Big] \,\mathrm{d}t \\
&= \tau \int_0^\tau (1 - \tfrac{t}{\tau}) \Big[- \tfrac{4}{\tau}(g_{y_k}(v_{k+1}, \psi_k) - g_{y_{k-1}}(v_k, \psi_k)) + 2 D_y g_{y_k}(\psi_k)(v_{k+1}, v_{k+1}) \\
&\qquad + \int_0^1 \Big(2\mathcal{W}_{,222}[y_{k-1}, y_{k-1} + r t v_k](v_k, \tfrac{1}{\tau}\psi_k, t v_k) \\
&\qquad\qquad - 2\mathcal{W}_{,222}[y_k, y_k + r t v_{k+1}](v_{k+1}, \tfrac{1}{\tau}\psi_k, t v_{k+1}) \\
&\qquad\qquad + \mathcal{W}_{,2212}[y_k, y_k + r t v_{k+1}](v_{k+1}, v_{k+1}, \psi_k, t v_{k+1}) \\
&\qquad\qquad + \mathcal{W}_{,2222}[y_k, y_k + r t v_{k+1}](v_{k+1}, v_{k+1}, \psi_k, t v_{k+1}) \Big) \,\mathrm{d}r \\
&\qquad + \mathcal{W}_{,222}[y_{k-1}, y_{k-1} + t v_k](v_k, v_k, \tfrac{t}{\tau}\psi_k) - \mathcal{W}_{,222}[y_k, y_k + t v_{k+1}](v_{k+1}, v_{k+1}, \tfrac{t}{\tau}\psi_k) \Big] \,\mathrm{d}t \\
&= -\tau^2 \left[2 \frac{g_{y_k}(v_{k+1}, \psi_k) - g_{y_{k-1}}(v_k, \psi_k)}{\tau} - D_y g_{y_k}(\psi_k)(v_{k+1}, v_{k+1}) \right] \\
&\quad + O\big[(\tau(\|v_k\|_{\mathbf{V}}^3 + \|v_{k+1}\|_{\mathbf{V}}^3) + (\tau\|v_k\|_{\mathbf{V}}^2 + \|v_{k+1} - v_k\|_{\mathbf{V}})(\|v_{k+1}\|_{\mathbf{V}} + \|v_k\|_{\mathbf{V}}))\tau^2 \|\psi_k\|_{\mathbf{V}}\big]
\end{aligned} \tag{6.6}$$

assuming sufficient differentiability of $\mathcal{W}$ and using the simple Taylor expansion formula

$$\begin{aligned}f[a_2,b_2](c_2,c_2,d)-f[a_1,b_1](c_1,c_1,d) &= f[a_1,b_1](c_2-c_1,c_2,d)+f[a_1,b_1](c_1,c_2-c_1,d)\\ &\quad + f_{,1}[a,b](c_2,c_2,d,a_2-a_1)+f_{,2}[a,b](c_2,c_2,d,b_2-b_1)\end{aligned}$$

for a spatially dependent, differentiable trilinear form f and $a=(1-\xi)a_1+\xi a_2$, $b=(1-\xi)b_1+\xi b_2$ for some $\xi\in[0,1]$.

Next, let us abbreviate $t_k=k\tau$. The smooth geodesic y satisfies

$$\begin{aligned}0&=-2\tfrac{\mathrm{d}}{\mathrm{d}t}g_{y(t)}(\dot y(t),\psi_k)|_{t=t_k}+D_yg_{y(t_k)}(\psi_k)(\dot y(t_k),\dot y(t_k))\\ &=-2\frac{g_{y(t_k)}(u_{k+1},\psi_k)-g_{y(t_{k-1})}(u_k,\psi_k)}{\tau}+D_yg_{y(t_k)}(\psi_k)(u_{k+1},u_{k+1})+O(\tau\|\psi_k\|_{\mathbf{V}})\end{aligned}$$

for $u_k=\frac{y(t_k)-y(t_{k-1})}{\tau}$ and $|O(\tau\|\psi_k\|_{\mathbf{V}})|\le C\tau(\|\dddot y\|_{\mathbf{V}}+\|\ddot y\|_{\mathbf{V}}(\|\dot y\|_{\mathbf{V}}+\tau\|\ddot y\|_{\mathbf{V}}))\|\psi_k\|_{\mathbf{V}}$. Now assume $y_{k+1}:=(\mathrm{EXP}^{k+1})_{y(0)}(\dot y(0)/K)$ exists for some $k<K$ (and thus also $y_0,\ldots,y_k$), then the left-hand side of (6.6) is zero. Subtracting $1/\tau^2$ times (6.6) from the above equation yields

$$\begin{aligned}&O\left[(\tau+\tau\|v_k\|_{\mathbf{V}}^3+\tau\|v_{k+1}\|_{\mathbf{V}}^3+(\tau\|v_k\|_{\mathbf{V}}^2+\|v_{k+1}-v_k\|_{\mathbf{V}})(\|v_{k+1}\|_{\mathbf{V}}+\|v_k\|_{\mathbf{V}}))\|\psi_k\|_{\mathbf{V}}\right]\\ &=2\frac{g_{y_k}(v_{k+1},\psi_k)-g_{y_{k-1}}(v_k,\psi_k)}{\tau}-2\frac{g_{y(t_k)}(u_{k+1},\psi_k)-g_{y(t_{k-1})}(u_k,\psi_k)}{\tau}\\ &\quad-D_yg_{y_k}(\psi_k)(v_{k+1},v_{k+1})+D_yg_{y(t_k)}(\psi_k)(u_{k+1},u_{k+1})\end{aligned}\tag{6.7}$$

Introducing $e_k=y_k-y(t_k)$ as well as $e_k^v=\frac{e_k-e_{k-1}}{\tau}$, the first line on the right-hand side can be rewritten as

$$\begin{aligned}&2\frac{g_{y_k}(v_{k+1}-v_k,\psi_k)-g_{y(t_k)}(u_{k+1}-u_k,\psi_k)}{\tau}\\ &\qquad+2D_yg_{y_k}(v_k)(v_k,\psi_k)-2D_yg_{y(t_k)}(u_k)(u_k,\psi_k)+O(\tau(\|v_k\|_{\mathbf{V}}^3+\|u_k\|_{\mathbf{V}}^3)\|\psi_k\|_{\mathbf{V}})\\ &=2g_{y_k}\left(\frac{e_{k+1}^v-e_k^v}{\tau},\psi_k\right)+O\big[(\|e_k\|_{\mathbf{V}}\|\tfrac{u_{k+1}-u_k}{\tau}\|_{\mathbf{V}}+\|e_k\|_{\mathbf{V}}\|u_k\|_{\mathbf{V}}^2\\ &\qquad+\|e_k^v\|_{\mathbf{V}}(\|u_k\|_{\mathbf{V}}+\|v_k\|_{\mathbf{V}})+\tau(\|v_k\|_{\mathbf{V}}^3+\|u_k\|_{\mathbf{V}}^3))\|\psi_k\|_{\mathbf{V}}\big]\,,\end{aligned}$$

and the second line is bounded by $O[(\|e_k\|_{\mathbf{V}}\|u_{k+1}\|_{\mathbf{V}}^2+\|e_{k+1}^v\|_{\mathbf{V}}\|u_{k+1}+v_{k+1}\|_{\mathbf{V}})\|\psi_k\|_{\mathbf{V}}]$. Thus, exploiting the boundedness of y and its derivatives as well as

$$\begin{aligned}&u_k=\dot y(t_k)+\tau O(\ddot y)\,,\quad u_{k+1}-u_k=\tau\ddot y(t_k)+\tau^2O(\dddot y)\,,\\ &\|v_k\|_{\mathbf{V}}\le\|u_k\|_{\mathbf{V}}+\|e_k^v\|_{\mathbf{V}}\,,\quad \|v_{k+1}-v_k\|_{\mathbf{V}}\le\|u_{k+1}-u_k\|_{\mathbf{V}}+\|e_{k+1}^v\|_{\mathbf{V}}+\|e_k^v\|_{\mathbf{V}}\,,\end{aligned}$$

(6.7) can be rewritten as

$$\begin{pmatrix}\frac{e_{k+1}-e_k}{\tau}\\ \frac{e_{k+1}^v-e_k^v}{\tau}\end{pmatrix}=\begin{pmatrix}e_{k+1}^v\\ O[\tau+\|e_k\|_{\mathbf{V}}+(\|e_k^v\|_{\mathbf{V}}+\|e_{k+1}^v\|_{\mathbf{V}})(1+\|e_k^v\|_{\mathbf{V}}+\|e_{k+1}^v\|_{\mathbf{V}})^2]\end{pmatrix}\,,$$

where we have also used the uniform coerciveness of g. If $\|e_j^v\|_{\mathbf{V}}\le 1$ for $j=0,\ldots,k+1$, then the above implies the existence of a constant $C>0$ such that with $U=C\left(\begin{smallmatrix}1&1\\1&1\end{smallmatrix}\right)$,

$$(\mathbb{1}-\tau U)\begin{pmatrix}\|e_{j+1}\|_{\mathbf{V}}\\ \|e_{j+1}^v\|_{\mathbf{V}}\end{pmatrix}\le(\mathbb{1}+\tau U)\begin{pmatrix}\|e_j\|_{\mathbf{V}}\\ \|e_j^v\|_{\mathbf{V}}\end{pmatrix}+C\tau^2\,,\qquad j=0,\ldots,k,$$

which together with $e_0 = 0$, $\|e_1^v\|_{\mathbf{V}} = O(\tau)$ by classical arguments implies $\|e_j\|_{\mathbf{V}}, \|e_j^v\|_{\mathbf{V}} \leq \kappa\tau$, $j = 0, \ldots, k+1$, for some $\kappa > 0$ which is independent of k and K.

Now choose $\frac{1}{\tau} = K$ large enough such that $C_K := \kappa\tau^2 + \tau \max_{t\in[0,1]} \|\dot{y}(t)\|_{\mathbf{V}} < \epsilon$ for $\epsilon \ll 1$ from Lemma 6.5. Then by induction, $(\mathrm{EXP}^k)(\dot{y}(0)/K) = y_k$ exists with $\|e_k\|_{\mathbf{V}}, \|e_k^v\|_{\mathbf{V}} < \kappa\tau$. Indeed, for $k = 0$ the situation is clear. Now if y_j exists with $\|e_j\|_{\mathbf{V}}, \|e_j^v\|_{\mathbf{V}} < \kappa\tau$ for all $j < k$, then $\|y_{k-1} - y_{k-2}\|_{\mathbf{V}} \leq \left(\|u_{k-1}\|_{\mathbf{V}} + \|e_{k-1}^v\|_{\mathbf{V}}\right)\tau \leq \left(\max_{t\in[0,1]} \|\dot{y}(t)\|_{\mathbf{V}} + \kappa\tau\right)\tau \leq C_K < \epsilon$ so that $y_k = (\mathrm{EXP}^2)_{y_{k-2}}(y_{k-1} - y_{k-2})$ exists by Lemma 6.5 with $y_k - y_{k-1} = y_{k-1} - y_{k-2} + O(\epsilon^{3/2})$. Thus $\|e_k\|_{\mathbf{V}}, \|e_k^v\|_{\mathbf{V}} < 1$, which via the above error estimates in turn implies $\|e_k\|_{\mathbf{V}}, \|e_k^v\|_{\mathbf{V}} < \kappa\tau$. □

THEOREM 6.10 (Existence and convergence of $\mathbf{P}_{y_K,\ldots,y_0}$). *Let $y : [0,1] \to \mathcal{M}$ be a smooth path, and let $\zeta : [0,1] \to \mathbf{V}$ be a parallel vector field along y. For $K \in \mathbb{N}$ and $\tau = \frac{1}{K}$, let $y_k = y(k\tau)$, then under the hypotheses (H1) and (H2), the discrete parallel transport fulfills $\|K\mathbf{P}_{y_K,\ldots,y_0}(\frac{\zeta(0)}{K}) - \zeta(1)\|_{\mathbf{V}} = O(\tau)$.*

Proof. At first, we examine the residual resulting from the evaluation of the discrete counterpart of the parallel transport equation on the interpolated continuous vector field ζ. ζ is defined to be parallel if $\frac{D}{\mathrm{d}t}\zeta = 0$ holds for the covariant derivative $\frac{D}{\mathrm{d}t}$ of ζ along the curve x, where the covariant derivative is defined via

$$g_{y(t)}((\tfrac{D}{\mathrm{d}t}\zeta)(t), \psi) = g_{y(t)}(\dot{\zeta}(t), \psi) + g_{y(t)}(\Gamma(\dot{y}(t), \zeta(t)), \psi) = 0 \tag{6.8}$$

for all test vectors $\psi \in \mathbf{V}$ and $t \in [0,1]$. Here the Christoffel tensor $\Gamma : \mathbf{V} \times \mathbf{V} \to \mathbf{V}$ is defined by $g_y(\Gamma(v,w), \psi) = \frac{1}{2}\left[(D_y g_y)(w)(\psi, v) + (D_y g_y)(v)(\psi, w) - (D_y g_y)(\psi)(v, w)\right]$ for all $v, w, \psi \in \mathbf{V}$. Let us abbreviate $t_k = k\tau$, $\zeta_k = \zeta(t_k)$, and $v_k = \frac{y_k - y_{k-1}}{\tau}$. Applying $g_y(v,w) = \frac{1}{2}\mathcal{W}_{,22}[y,y](v,w)$ (Lemma 5.7) and (5.10) as well as the approximations $\dot{\zeta}(t_{k-1}) = \frac{\zeta_k - \zeta_{k-1}}{\tau} + O(\tau\|\ddot{\zeta}\|_{\mathbf{V}})$, $\dot{y}(t_{k-1}) = v_k + O(\tau\|\ddot{y}\|_{\mathbf{V}})$, (6.8) turns into

$$\begin{aligned}
&O\Big[\tau\big(\sup_{t\in[0,1]}\|\ddot{\zeta}(t)\|_{\mathbf{V}} + \sup_{t\in[0,1]}\|\ddot{y}(t)\|_{\mathbf{V}} \sup_{t\in[0,1]}\|\zeta(t)\|_{\mathbf{V}}\big)\|\psi\|_{\mathbf{V}}\Big]\\
&= \mathcal{W}_{,22}[y_{k-1}, y_{k-1}](\tfrac{\zeta_k - \zeta_{k-1}}{\tau}, \psi)\\
&\quad + \tfrac{1}{2}\Big(\mathcal{W}_{,221}[y_{k-1}, y_{k-1}](\psi, v_k, \zeta_{k-1}) + \mathcal{W}_{,222}[y_{k-1}, y_{k-1}](\psi, v_k, \zeta_{k-1})\\
&\qquad + \mathcal{W}_{,221}[y_{k-1}, y_{k-1}](\psi, \zeta_{k-1}, v_k) + \mathcal{W}_{,222}[y_{k-1}, y_{k-1}](\psi, \zeta_{k-1}, v_k)\\
&\qquad - \mathcal{W}_{,221}[y_{k-1}, y_{k-1}](v_k, \zeta_{k-1}, \psi) - \mathcal{W}_{,222}[y_{k-1}, y_{k-1}](v_k, \zeta_{k-1}, \psi)\Big)\\
&= \mathcal{W}_{,22}[y_{k-1}, y_{k-1}](\tfrac{\zeta_k - \zeta_{k-1}}{\tau}, \psi)\\
&\quad + \tfrac{1}{2}\Big(- \mathcal{W}_{,211}[y_{k-1}, y_{k-1}](\psi, v_k, \zeta_{k-1}) - \mathcal{W}_{,212}[y_{k-1}, y_{k-1}](\psi, v_k, \zeta_{k-1})\\
&\qquad + \mathcal{W}_{,221}[y_{k-1}, y_{k-1}](\psi, \zeta_{k-1}, v_k) - \mathcal{W}_{,221}[y_{k-1}, y_{k-1}](v_k, \zeta_{k-1}, \psi)\Big)\\
&= \mathcal{W}_{,22}[y_{k-1}, y_{k-1}](\tfrac{\zeta_k - \zeta_{k-1}}{\tau}, \psi)\\
&\quad - \tfrac{1}{2}\Big(\mathcal{W}_{,211}[y_{k-1}, y_{k-1}](\psi, v_k, \zeta_{k-1}) + \mathcal{W}_{,221}[y_{k-1}, y_{k-1}](v_k, \zeta_{k-1}, \psi)\Big)\\
&= \mathcal{W}_{,22}[y_{k-1}, y_{k-1}](\tfrac{\zeta_k - \zeta_{k-1}}{\tau}, \psi) - \frac{\mathcal{W}_{,112}[y_{k-1}, y_{k-1}] + \mathcal{W}_{,221}[y_{k-1}, y_{k-1}](\zeta_{k-1}, v_k, \psi)}{2},
\end{aligned}$$

where we have used Schwarz's theorem and the identity

$$\begin{aligned}
\mathcal{W}_{,221}[y,y] + \mathcal{W}_{,222}[y,y] &= \mathcal{W}_{,111}[y,y] + \mathcal{W}_{,112}[y,y]\\
&= -\mathcal{W}_{,121}[y,y] - \mathcal{W}_{,122}[y,y] = -\mathcal{W}_{,211}[y,y] - \mathcal{W}_{,212}[y,y],
\end{aligned}$$

which can be derived by differentiating the identity $\mathcal{W}_{,22}[y,y] = \mathcal{W}_{,11}[y,y] = -\mathcal{W}_{,12}[y,y] = -\mathcal{W}_{,21}[y,y]$ from Lemma 5.7 with respect to y. Due to the smoothness of ζ and y we achieve the estimate

$$\begin{aligned}&\mathcal{W}_{,22}[y_{k-1},y_{k-1}](\tfrac{\zeta_k-\zeta_{k-1}}{\tau},\psi)\\ &-\tfrac{1}{2}\Big(\mathcal{W}_{,112}[y_{k-1},y_{k-1}]+\mathcal{W}_{,221}[y_{k-1},y_{k-1}]\Big)(\zeta_{k-1},v_k,\psi) = O(\tau\|\psi\|_{\mathbf{V}})\end{aligned} \tag{6.9}$$

Next, we derive an estimate for the residual obtained when evaluating the discrete counterpart of the parallel transport equation on the discrete solution. Thus, let us abbreviate $\xi_k = \mathbf{P}_{y_k,\dots,y_0}(\zeta(0)/K)$ and denote the center of the k^{th} parallelogram in Schild's ladder by y_k^c. ξ_k and y_k^c satisfy for all $\psi \in \mathbf{V}$ the two nonlinear equations

$$\mathcal{W}_{,2}[y_{k-1}+\xi_{k-1},y_k^c](\psi)+\mathcal{W}_{,1}[y_k^c,y_k](\psi) = 0\,, \tag{6.10}$$

$$\mathcal{W}_{,2}[y_{k-1},y_k^c](\psi)+\mathcal{W}_{,1}[y_k^c,y_k+\xi_k](\psi) = 0\,, \tag{6.11}$$

which for the moment we assume to be uniquely solvable. Upon Taylor expansion about (y_k^c,y_k^c) analogously to (5.9), one obtains

$$\begin{aligned}\mathcal{W}[y_{k-1}+\xi_{k-1},y_k^c] &= \int_0^1 (1-s)\mathcal{W}_{,11}[y_k^c+s(y_{k-1}+\xi_{k-1}-y_k^c),y_k^c]\\ &\qquad\qquad (y_{k-1}+\xi_{k-1}-y_k^c,y_{k-1}+\xi_{k-1}-y_k^c)\,\mathrm{d}s\,,\\ \mathcal{W}[y_k^c,y_k] &= \int_0^1 (1-s)\mathcal{W}_{,22}[y_k^c,y_k^c+s(y_k-y_k^c)](y_k-y_k^c,y_k-y_k^c)\,\mathrm{d}s\,.\end{aligned}$$

Differentiating these expressions, the first equation (6.10) defining the discrete parallel transport turns into

$$\begin{aligned}0 = \int_0^1 (1-s)\Big[&-2\mathcal{W}_{,11}[y_k^c,y_k^c](\psi,y_{k-1}+\xi_{k-1}-y_k^c)\\ &\quad-2\int_0^1 \mathcal{W}_{,111}[y_k^c+rs(y_{k-1}+\xi_{k-1}-y_k^c),y_k^c](\psi,y_{k-1}+\xi_{k-1}-y_k^c,s(y_{k-1}+\xi_{k-1}-y_k^c))\,\mathrm{d}r\\ &+\mathcal{W}_{,111}[y_k^c+s(y_{k-1}+\xi_{k-1}-y_k^c),y_k^c](y_{k-1}+\xi_{k-1}-y_k^c,y_{k-1}+\xi_{k-1}-y_k^c,(1-s)\psi)\\ &+\mathcal{W}_{,112}[y_k^c+s(y_{k-1}+\xi_{k-1}-y_k^c),y_k^c](y_{k-1}+\xi_{k-1}-y_k^c,y_{k-1}+\xi_{k-1}-y_k^c,\psi)\\ &-2\mathcal{W}_{,22}[y_k^c,y_k^c](\psi,y_k-y_k^c)\\ &\quad-2\int_0^1 \mathcal{W}_{,222}[y_k^c,y_k^c+rs(y_k-y_k^c)](\psi,y_k-y_k^c,s(y_k-y_k^c))\,\mathrm{d}r\\ &+\mathcal{W}_{,221}[y_k^c,y_k^c+s(y_k-y_k^c)](y_k-y_k^c,y_k-y_k^c,\psi)\\ &+\mathcal{W}_{,222}[y_k^c,y_k^c+s(y_k-y_k^c)](y_k-y_k^c,y_k-y_k^c,(1-s)\psi)\Big]\,\mathrm{d}s\\ = -\mathcal{W}_{,11}[y_k^c,&y_k^c](\psi,y_{k-1}+\xi_{k-1}-y_k^c)-\mathcal{W}_{,22}[y_k^c,y_k^c](\psi,y_k-y_k^c)\\ +\tfrac{1}{2}\mathcal{W}_{,112}&[y_k^c,y_k^c](y_{k-1}+\xi_{k-1}-y_k^c,y_{k-1}+\xi_{k-1}-y_k^c,\psi)\\ +\tfrac{1}{2}\mathcal{W}_{,221}&[y_k^c,y_k^c](y_k-y_k^c,y_k-y_k^c,\psi)\\ +O\big[(\|y_{k-1}&+\xi_{k-1}-y_k^c\|_{\mathbf{V}}^3+\|y_k-y_k^c\|_{\mathbf{V}}^3)\|\psi\|_{\mathbf{V}}\big]\,.\end{aligned}$$

Here we used that for a spatially differentiable trilinear form f the relation $-2f[a,a+rsv](s\psi,v,v)+f[a,a+sv]((1-s)\psi,v,v) = f[a,a]((1-3s)\psi,v,v)+O(\|v\|^3\|\psi\|)$

holds for $0 \leq r, s \leq 1$, which after multiplication with $(1-s)$ integrates up to $O(\|v\|^3\|\psi\|)$. Likewise, the second equation (6.11) turns into

$$\begin{aligned}0 = &-\mathcal{W}_{,11}[y_k^c, y_k^c](\psi, y_{k-1} - y_k^c) - \mathcal{W}_{,22}[y_k^c, y_k^c](\psi, y_k + \xi_k - y_k^c)\\ &+\tfrac{1}{2}\mathcal{W}_{,112}[y_k^c, y_k^c](y_{k-1} - y_k^c, y_{k-1} - y_k^c, \psi)\\ &+\tfrac{1}{2}\mathcal{W}_{,221}[y_k^c, y_k^c](y_k + \xi_k - y_k^c, y_k + \xi_k - y_k^c, \psi)\\ &+O\left[(\|y_{k-1} - y_k^c\|_{\mathbf{V}}^3 + \|y_k + \xi_k - y_k^c\|_{\mathbf{V}}^3)\|\psi\|_{\mathbf{V}}\right]\end{aligned}$$

Subtracting the second from the first equation, dividing by τ^2, and using $\mathcal{W}_{,11} = \mathcal{W}_{,22}$, one obtains

$$\begin{aligned}0 = &\,\mathcal{W}_{,22}[y_k^c, y_k^c](\tfrac{\frac{1}{\tau}\xi_k - \frac{1}{\tau}\xi_{k-1}}{\tau}, \psi)\\ &+\tfrac{1}{2}\Big[\mathcal{W}_{,112}[y_k^c, y_k^c](\tfrac{\xi_{k-1}}{\tau}, \tfrac{\xi_{k-1}+2(y_{k-1}-y_k^c)}{\tau}, \psi) - \mathcal{W}_{,221}[y_k^c, y_k^c](\tfrac{\xi_k}{\tau}, \tfrac{\xi_k+2(y_k-y_k^c)}{\tau}, \psi)\Big]\\ &+O\left[(\|y_{k-1} - y_k^c\|_{\mathbf{V}}^3 + \|y_k - y_k^c\|_{\mathbf{V}}^3 + \|\xi_{k-1}\|_{\mathbf{V}}^3 + \|\xi_k\|_{\mathbf{V}}^3)\|\psi\|_{\mathbf{V}}\right]\,. \end{aligned} \tag{6.12}$$

Finally, we derive an equation for the error propagation and subtract (6.9) from (6.12). Introducing the error $e_k = K\xi_k - \zeta_k = \frac{\xi_k}{\tau} - \zeta_k$, we arrive at

$$\begin{aligned}&\mathcal{W}_{,22}[y_k^c, y_k^c](\tfrac{e_k - e_{k-1}}{\tau}, \psi) = \tfrac{1}{2}\big(\mathcal{W}_{,112}[y_k^c, y_k^c] + \mathcal{W}_{,221}[y_k^c, y_k^c]\big)(e_{k-1}, v_k, \psi)\\ &+O\Big[\big[\tau + \|y_{k-1} - y_k^c\|_{\mathbf{V}}^3 + \|y_k - y_k^c\|_{\mathbf{V}}^3 + \|\xi_{k-1}\|_{\mathbf{V}}^3 + \|\xi_k\|_{\mathbf{V}}^3 + \|y_k^c - y_{k-1}\|_{\mathbf{V}}\|\tfrac{\zeta_k - \zeta_{k-1}}{\tau}\|_{\mathbf{V}}\\ &+\|y_k^c - y_{k-1}\|_{\mathbf{V}}\|\zeta_{k-1}\|_{\mathbf{V}}\|v_k\|_{\mathbf{V}} + \tfrac{1}{\tau^2}\|\xi_{k-1}\|_{\mathbf{V}}\|\xi_{k-1} - 2(y_k^c - \tfrac{y_k + y_{k-1}}{2})\|_{\mathbf{V}}\\ &+\tfrac{1}{\tau}\|\xi_k - \xi_{k-1}\|_{\mathbf{V}}\|v_k\|_{\mathbf{V}} + \tfrac{1}{\tau^2}\|\xi_k\|_{\mathbf{V}}\|\xi_k - 2(y_k^c - \tfrac{y_k + y_{k-1}}{2})\|_{\mathbf{V}}\big]\|\psi\|_{\mathbf{V}}\Big]\,.\end{aligned} \tag{6.13}$$

Applying the boundedness of v_k, ζ_k, $\dot{\zeta}$, and the uniform estimates

$$\begin{aligned}\|\tfrac{\xi_k}{\tau} - \tfrac{\xi_{k-1}}{\tau}\|_{\mathbf{V}} &\leq \|\zeta_k - \zeta_{k-1}\|_{\mathbf{V}} + \|e_k\|_{\mathbf{V}} + \|e_{k-1}\|_{\mathbf{V}}\\ &= O(\tau + \|e_k\|_{\mathbf{V}} + \|e_{k-1}\|_{\mathbf{V}})\\ \|\tfrac{\xi_k}{\tau}\|_{\mathbf{V}} &\leq \|\zeta_k\|_{\mathbf{V}} + \|e_k\|_{\mathbf{V}}\\ \|y_k^c - y_k\|_{\mathbf{V}} &= O(\|y_{k-1} + \xi_{k-1} - y_k\|_{\mathbf{V}})\\ &= \tau O(\|v_k\|_K + \|\zeta_{k-1}\|_{\mathbf{V}} + \|e_{k-1}\|_{\mathbf{V}})\\ \|y_k^c - y_{k-1}\|_{\mathbf{V}} &= O(\|y_k + \xi_k - y_{k-1}\|_{\mathbf{V}})\\ &= \tau O(\|v_k\|_{\mathbf{V}} + \|\zeta_k\|_{\mathbf{V}} + \|e_k\|_{\mathbf{V}})\\ \|\xi_{k-1} - 2(y_k^c - \tfrac{y_k + y_{k-1}}{2})\|_{\mathbf{V}} &= O(\|y_{k-1} + \xi_{k-1} - y_k\|_{\mathbf{V}}^2)\\ &= \tau^2 O((\|v_k\|_{\mathbf{V}} + \|\zeta_{k-1}\|_{\mathbf{V}} + \|e_{k-1}\|_{\mathbf{V}})^2)\\ \|\xi_k - 2(y_k^c - \tfrac{y_k + y_{k-1}}{2})\|_{\mathbf{V}} &= O(\|y_{k-1} - y_k - \xi_k\|_{\mathbf{V}}^2)\\ &= \tau^2 O((\|v_k\|_K + \|\zeta_k\|_{\mathbf{V}} + \|e_k\|_{\mathbf{V}})^2)\end{aligned}$$

(the last four follow from Remark 6.8, assuming $\|\xi_{k-1}\|_{\mathbf{V}}, \|\xi_k\|_{\mathbf{V}}$ to be sufficiently small), we find

$$\tfrac{e_k - e_{k-1}}{\tau} = O[\|e_{k-1}\|_{\mathbf{V}} + \tau(1 + \|e_{k-1}\|_{\mathbf{V}})^3 + \|e_k\|_{\mathbf{V}} + \tau(1 + \|e_k\|_{\mathbf{V}})^3]\,. \tag{6.14}$$

As in the proof of Theorem 6.9, this implies the existence of a constant $C > 0$ (independent of K) such that if $\|e_j\|_{\mathbf{V}} \leq 1$ for $j \leq k \leq K$, then $\|e_k\|_{\mathbf{V}} \leq C\tau$.

We now inductively show the existence of ξ_k with $\|e_k\|_{\mathbf{V}} \leq C\tau$. Choose K large enough such that $C+\sup_{k=1,\ldots,K} \|\zeta_{k-1}\|_{\mathbf{V}} + \|v_k\|_{\mathbf{V}} + \|\zeta_k\|_{\mathbf{V}} < \epsilon K$ for a given $\epsilon \ll 1$ from Lemma 6.5. Clearly, ξ_0 exists, and $e_0 = 0$. Now assume the existence of ξ_j, $j < k$, with $\|e_j\|_{\mathbf{V}} \leq C\tau$. In particular, this implies $\|\xi_{k-1}\|_{\mathbf{V}} + \tau\|v_k\|_{\mathbf{V}} < \epsilon$ and thus by Lemmata 6.2 and 6.5 the existence of y_k^c and ξ_k. Furthermore, Remark 6.8 ensures $\|e_k\|_{\mathbf{V}} \leq 1$ such that the above estimates imply $\|e_k\|_{\mathbf{V}} \leq C\tau$. ◻

The parallel transport also converges if the interpolated smooth path $(y_0, \ldots, y_K)$ is replaced by another approximating path, for instance a discrete geodesic.

COROLLARY 6.11. *Let $y : [0,1] \to \mathcal{M}$ be a smooth path and $\zeta : [0,1] \to \mathbf{V}$ a parallel vector field along y. For $K \in \mathbb{N}$ and $\tau = \frac{1}{K}$, let y_k satisfy $\|y_k - y(k\tau)\|_{\mathbf{V}} \leq \epsilon$, $k = 0, \ldots, K$, then under the hypotheses (H1) and (H2), $\|K\mathbf{P}_{y_K,\ldots,y_0}(\frac{\zeta(0)}{K}) - \zeta(1)\|_{\mathbf{V}} = O(\tau + \epsilon)$.*

Proof. Following the previous proof we arrive at a slightly altered version of (6.13),

$$\begin{aligned}
&\mathcal{W}_{,22}[y_k^c, y_k^c](\tfrac{e_k - e_{k-1}}{\tau}, \psi) = \tfrac{1}{2}\big(\mathcal{W}_{,112}[y_k^c, y_k^c] + \mathcal{W}_{,221}[y_k^c, y_k^c]\big)(e_{k-1}, v_k, \psi) \\
&+ O\Big[\big[\tau + \|y_{k-1} - y_k^c\|_{\mathbf{V}}^3 + \|y_k - y_k^c\|_{\mathbf{V}}^3 + \|\xi_{k-1}\|_{\mathbf{V}}^3 + \|\xi_k\|_{\mathbf{V}}^3 + \|y_k^c - y(t_{k-1})\|_{\mathbf{V}} \|\tfrac{\zeta_k - \zeta_{k-1}}{\tau}\|_{\mathbf{V}} \\
&+ \|y_k^c - y(t_{k-1})\|_{\mathbf{V}} \|\zeta_{k-1}\|_{\mathbf{V}} \|v_k\|_{\mathbf{V}} + \tfrac{1}{\tau^2}\|\xi_{k-1}\|_{\mathbf{V}} \|\xi_{k-1} - 2(y_k^c - \tfrac{y_k + y_{k-1}}{2})\|_{\mathbf{V}} \\
&+ \tfrac{1}{\tau}\|\xi_k - \xi_{k-1}\|_{\mathbf{V}} \|v_k\|_{\mathbf{V}} + \tfrac{1}{\tau^2}\|\xi_k\|_{\mathbf{V}} \|\xi_k - 2(y_k^c - \tfrac{y_k + y_{k-1}}{2})\|_{\mathbf{V}}\big] \|\psi\|_{\mathbf{V}}\Big],
\end{aligned}$$

where $t_k = k\tau$ and $v_k = \frac{y(t_k) - y(t_{k-1})}{\tau}$. Using $\|y_k^c - y(t_{k-1})\|_{\mathbf{V}} \leq \|y_k^c - y_{k-1}\|_{\mathbf{V}} + \epsilon$ as well as the estimates from the previous proof we obtain (6.14) with $O(\epsilon)$ added to the right-hand side, from which the claim follows by the same arguments. ◻

COROLLARY 6.12. *Let $\theta \in \mathbf{V}$ and $\eta : y_A + \mathbf{V} \to \mathbf{V}$ be a smooth vector field. Define $\eta^\tau = (\eta(y), \eta(y + \tau\theta))$, then under the hypotheses (H1) and (H2) we have $\|\frac{1}{\tau^2}\boldsymbol{\nabla}_{\tau\theta}(\tau\eta^\tau) - \nabla_\theta \eta\|_{\mathbf{V}} = O(\tau)$.*

Proof. Consider the continuous geodesic $y(t)$ with $y(0) = y$ and $\dot{y}(0) = \theta$. We have $\|y(\tau) - (y + \tau\theta)\|_{\mathbf{V}} = O(\tau^2)$. Now define $\xi_1 = \tau\eta(y + \tau\theta)$ and $\xi_0 = \mathbf{P}_{y+\tau\theta,y}^{-1}\xi_1$ as well as $\zeta_1 = \eta(y(\tau))$ and ζ_0 as the vector ζ_1 parallel transported from $y(\tau)$ to $y(0)$. Furthermore introduce the error $e_k = \frac{1}{\tau}\xi_k - \zeta_k$, $k = 0, 1$. Since $\mathbf{P}_{y+\tau\theta,y}^{-1}$ is defined via the same discrete Euler–Lagrange equations ((2.6) and (2.7) for $k = 0$, $y_0 = y$, and $y_1 = y + \tau\theta$) as $\mathbf{P}_{y+\tau\theta,y}$, (6.14) holds also here, only with $O(\|y(\tau) - (y + \tau\theta)\|_{\mathbf{V}}) = O(\tau^2)$ added to the right-hand side (exactly as in the previous corollary). Using $e_1 = O(\tau^2)$, we immediately see $e_0 = O(\tau^2)$. The relation $\nabla_\theta \eta = \frac{\zeta_0 - \eta(y)}{\tau} + O(\tau)$ together with $\zeta_0 + O(\tau^2) = \frac{1}{\tau}\xi_0 = \eta(y) + \frac{1}{\tau}\boldsymbol{\nabla}_{\tau\theta}(\tau\eta^\tau)$ now implies the claim. ◻

7. Application to shape spaces.

In this section we will discuss the scope of applications of the presented convergence theory. In general, it can be utilized in a straightforward fashion for finite-dimensional manifolds and for those infinite-dimensional manifolds where motion paths are compact in state space and the metric can be rephrased properly on Lagrangian velocity fields. In what follows we exemplify this in the case of the discrete geodesic calculus on embedded manifolds from Section 4, on the parametrized finite-dimensional shape space of viscous-fluid objects from Section 3.1, and on the simplified shape space of viscous one-dimensional shells from Section 3.2.

In its current form the convergence theory is not yet applicable to the flow of diffeomorphism approach in Section 3.3. The difficulty lies in identifying the manifold $\mathcal{M}$ locally with an affine space $y_A + \mathbf{V}$ of equivalent topology, in particular if shape variations are described in infinite dimensions by Eulerian instead of Lagrangian velocity fields.

7.1. On embedded finite-dimensional manifolds (*cf*. Section 4).

The most basic example has been given in Section 4 by a smooth, complete, $(m-1)$-dimensional manifold $\mathcal{M}$, embedded in $\mathbb{R}^m$. Due to the assumptions on $\mathcal{M}$, a continuous geodesic exists between any two points $y_A, y_B \in \mathcal{M}$. Furthermore, we assume the embedding to be such that (5.5) holds for the function $\mathcal{W}$ which represents the squared extrinsic distance between two points on $\mathcal{M}$.

Even though in this example $\mathcal{M}$ is not an affine space, the existence proof for discrete geodesics (Theorem 5.4) still applies if in it we replace $\mathbf{Y}$ by $\mathcal{M}$ (note that due to the finite dimensions, the proof no longer has to rely on a compact embedding of $\mathcal{M}$ into some larger space). All subsequent results either do not make use of the manifold structure or are essentially local in the sense that their proofs only consider a neighborhood of a continuous geodesic. This characteristic enables us to transfer the results onto the example at hand: Via a chart we locally identify the manifold $\mathcal{M}$ with an open subset of $\mathbb{R}^{m-1}$ and set $\mathbf{Y} = \mathbf{V} = \mathbb{R}^{m-1}$. In more detail, all convergence proofs involve a continuous geodesic or smooth path (the limit object of convergence), around which a small compact tubular neighborhood U can be identified with a compact subset $\phi(U)$ of $\mathbb{R}^{m-1}$ (where $\phi : U \to \mathbb{R}^{m-1}$ is the parametrization). The induced metric on $\phi(U)$ is clearly equivalent to the Euclidean metric. Hence, if we consider $\phi(U) \subset \mathbf{Y}$ instead of $\mathcal{M}$, we are in the setting of Sections 5 and 6, and all convergence results apply. Now, convergence of paths y_τ in $W^{1,2}((0,1); \phi(U))$ and of vectors $\frac{1}{\tau} v_\tau \in \mathbf{V}$ against vectors v in a tangent space $T_{\phi(y)}\phi(U)$ immediately implies the same rate convergence of geodesics $\phi^{-1}(y_\tau)$ in $W^{1,2}((0,1); U)$ and of $\frac{1}{\tau}(\phi^{-1}(\phi(y) + v_\tau) - y) \in \mathbb{R}^m$ against $D\phi^{-1} v \in T_y U$ so that discrete geodesics, logarithm, exponential map, and parallel transport converge as well in this sense. From the proofs it should be clear that the restriction to U does not impair the results.

7.2. On the parametrized finite-dimensional shape space of viscous-fluid objects (*cf.* Section 3.1). If we consider viscous fluid objects being represented by spline-parametrized boundary contours, each shape can be represented (uniquely up to rigid body motions) by an n-tuple $(q_1, \ldots, q_n)$ of control points in $\mathbb{R}^d$ ($d = 2$ or $d = 3$). First we quotient out the rigid body motion invariance. To this end we fix a reference configuration $y_A = (\hat{q}_1, \ldots, \hat{q}_n) \in \mathbb{R}^{nd}$ with $\sum_{k=1}^n \hat{q}_k = 0$ and only consider those shapes $y = (q_1, \ldots, q_n)$ that can be reached by a displacement of y_A with zero average linear and angular momentum, *i.e.* $y \in \mathcal{M} = y_A + \mathbf{V}$ for $\mathbf{V} = \{(\dot{q}_1, \ldots, \dot{q}_n) \in \mathbb{R}^{nd} \,|\, \sum_{k=1}^n \dot{q}_k = 0 \text{ and } \sum_{k=1}^n \hat{q}_k \wedge \dot{q}_k = 0\}$. Also, we define $\mathbf{Y} = \mathbf{V}$. Then $\mathbf{Y}$ is a k-dimensional subspace of $\mathbb{R}^{nd}$ ($k = 2(n-1) - 1$ for $d = 2$ and $k = 3(n-2)$ for $d = 3$), of which the admissible shapes actually only make up a closed subset, *i.e.* a manifold with boundary, since shapes with self-intersection of the shape contour are not allowed. In fact, we shall restrict the admissible shapes to those whose boundaries satisfy a uniform cone condition, *i.e.* each point on the shape boundary must be the tip of two open cones with fixed opening angle α and height r, one contained inside and one outside the shape. In this context, the metric is given by $g_y(\dot{y}, \dot{y}) = \int_{\mathcal{O}[y]} \frac{\lambda}{2}(\mathrm{tr}\epsilon[\tilde{v}])^2 + \mu \mathrm{tr}(\epsilon[\tilde{v}]^2)\,\mathrm{d}x$ for $y = (q_1, \ldots, q_n) \in \mathbf{Y}$, $\dot{y} = (\dot{q}_1, \ldots, \dot{q}_n) \in \mathbf{V}$, where $\mathcal{O}[y]$ is the interior of the spline described by y, and $\tilde{v} : \mathcal{O}[y] \to \mathbb{R}^d$ is the velocity field minimizing the right hand side with Dirichlet boundary data prescribed by the variation of y and thus of all points on $\partial\mathcal{O}[y]$. Using Korn's inequality, it is a classical result in the direct method of the calculus of variations that a minimizing $\tilde{v}$ exists. This $\tilde{v}$ depends linearly on $\dot{y}$, hence g_y is indeed a bounded quadratic functional. Again by Korn's inequality, $g_y(\dot{y}, \dot{y}) = 0$ implies $\tilde{v}(x) = v_0 + Sx$ for a constant velocity v_0 and a skew-symmetric matrix S. However, $v_0 \neq 0$ or $S \neq 0$ contradict $\dot{y} = (\tilde{v}(q_1), \ldots, \tilde{v}(q_n)) \in \mathbf{V}$, which implies that g_y is positive definite, and the shape space indeed carries the structure of a Riemannian manifold. Furthermore, by the existence and regularity theory of linearized elasticity [6], using the implicit function theorem, the metric g_y and the functional $\mathcal{W}$ actually smoothly depend on y with $g_y = \frac{1}{2}\mathcal{W}_{,22}[y, y]$ so that (2.4) holds by Lemma 5.7. Furthermore, by the direct method in the calculus of variations we see that for points y_A and y_B which are sufficiently close there exists an energy minimizing path (the imposed cone condition for the shapes renders the existence of smooth geodesic paths for arbitrary end shapes y_A and y_B unclear). Now, from the coercivity property (5.5), we know that any discrete K-path between two points $y_A, y_B \in \mathcal{M}$ with smaller energy than $\mathrm{E}[(y_A, \dots, y_A, y_B)]$ has to lie completely within a radius of $K\gamma^{-1}(\mathrm{E}[(y_A, \dots, y_A, y_B)]/K)$ around y_A. Hence, in the proof of Theorem 5.4 we may restrict to that compact region. This yields existence of a discrete K-geodesic for any K. Theorem 5.6 now implies that all K-geodesics between y_A and y_B stay inside the same closed bounded and thus compact region. Hence, during all our convergence analysis we may restrict to a compact subset of $\mathcal{M} = \mathbf{V} = \mathbf{Y}$ on which the Euclidean metric and g are equivalent, hence all convergence results hold.

7.3. On the shape space of viscous shells (*cf*. Section 3.2). For the simplified model of one-dimensional shells (3.3), (3.4) let us consider for $0 \leq \alpha < \frac{1}{2}$ the space $\mathbf{Y} = \{y \in C^{1,\alpha}(\mathbb{R};\mathbb{R}^2) \,|\, y(s+1) = y(s)\}$ of $C^{1,\alpha}$-smooth closed curves represented by periodic parametrizations. Furthermore, let $\mathbf{V} = \{v \in W^{2,2}(\mathbb{R};\mathbb{R}^2) \,|\, v(s+1) = v(s),\ \int_0^1 v(s)\,\mathrm{d}s = 0\,, \int_0^1 v(s) \wedge y_A(s)\,\mathrm{d}s = 0\}$ for some fixed $y_A \in \mathbf{Y}$ be the space of velocity fields with zero average speed and angular momentum. Furthermore, we assume that $|y_{A,s}| \geq c > 0$. Then, we identify the manifold $\mathcal{M}$ of one-dimensional shells locally with an open subset of $y_A + \mathbf{V}$. By Sobolev embedding results $\mathbf{V}$ is compactly embedded in $\mathbf{Y}$, and $\mathbf{V}$ is a reflexive, separable space. Given $v \in L^2((0,1);\mathbf{V})$ with $\int_0^1 \|v(t,\cdot)\|^2_{\mathcal{W}^{2,2}((0,1),\mathbb{R}^2)}\,\mathrm{d}t \leq \epsilon$ for sufficiently small ϵ we obtain from $y(t,s) = y_A(s) + \int_0^t v(r,s)\,\mathrm{d}r$ that $\frac{c}{2} \leq |y_{,s}(t,s)| \leq C$ for the path $y \in L^2((0,1);\mathcal{M})$ with $v = \dot{y}$. Furthermore, for g_y defined in (3.4) $g_y(v,v) = 0$ implies $v = 0$. From this we deduce that in a sufficiently small neighborhood of y_A the metric g is uniformly bounded and uniformly coercive on $\mathbf{V}$ in the sense of (5.1) and that $y \mapsto g_y$ and $\mathcal{W}$ defined in (3.3) are smooth. Furthermore, the coercivity assumption (5.5) for $\mathcal{W}$ is fulfilled. Hence, for $y_B - y_A$ sufficiently small the direct method in the calculus of variations used in the proofs of Theorem 5.3 and Theorem 5.4 can be applied to establish the existence of a continuous and a discrete geodesic. Furthermore, Theorem 5.6 implies that a K-geodesic between y_A and y_B stays inside the same bounded region in $y_A + \mathbf{V}$. As above, the convergence analysis can be restricted to such a small neighborhood of y_A.

For the discrete approach on triangular meshes of fixed topology we describe a shape via the vertex positions and choose $\mathbf{Y} = \mathbf{V} = \mathbb{R}^{nd}$, where n is the number of nodes. Due to the smoothness of the energy $\mathcal{W}$ (3.2) our convergence theory directly applies even though the geometric interpretation of the associated metric $g_y = \frac{1}{2}\mathcal{W}_{,22}[y,y]$ is still open.

Acknowledgements. The authors thank Behrend Heeren for providing Figure 3.4.

REFERENCES

[1] Fred Almgren, Jean E. Taylor, and Lihe Wang, *Curvature-driven flows: a variational approach*, SIAM J. Control Optim., 31 (1993), pp. 387–438. 2

[2] V. I. Arnold, *Mathematical methods of classical mechanics*, Springer-Verlag, New York, 1978. Translated from the Russian by K. Vogtmann and A. Weinstein, Graduate Texts in Mathematics, 60. 2

[3] M. F. Beg, M.I. Miller, A. Trouvé, and L. Younes, *Computational anatomy: Computing metrics on anatomical shapes*, in Proceedings of 2002 IEEE ISBI, 2002, pp. 341–344. 1, 9

[4] M. Faisal Beg, Michael I. Miller, Alain Trouvé, and Laurent Younes, *Computing large deformation metric mappings via geodesic flows of diffeomorphisms*, International Journal of Computer Vision, 61 (2005), pp. 139–157. 2

[5] Jean-David Benamou and Yann Brenier, *A computational fluid mechanics solution to the Monge-Kantorovich mass transfer problem*, Numer. Math., 84 (2000), pp. 375–393. 2

[6] Philippe G. Ciarlet, *Mathematical Elasticity, Vol. I: Three–Dimensional Elasticity*, Studies in Mathematics and its Applications, Elsevier, Amsterdam, 1997. 28

[7] D. Dupuis, U. Grenander, and M.I. Miller, *Variational problems on flows of diffeomorphisms for image matching*, Quarterly of Applied Mathematics, 56 (1998), pp. 587–600. 2, 9

[8] J. Ehlers, F. A. E. Pirani, and A. Schild, *The geometry of free fall and light propagation*, in General relativity (papers in honour of J. L. Synge), Clarendon Press, Oxford, 1972, pp. 63–84. 5

[9] P.T. Fletcher, Conglin Lu, S.M. Pizer, and Sarang Joshi, *Principal geodesic analysis for the study of nonlinear statistics of shape*, Medical Imaging, IEEE Transactions on, 23 (2004), pp. 995–1005. 1

[10] Gero Friesecke, Richard James, Maria Giovanna Mora, and Stefan Müller, *Derivation of nonlinear bending theory for shells from three-dimensional nonlinear elasticity by Gamma-convergence*, C. R. Math. Acad. Sci. Paris 336, 8 (2003), pp. 697–702. 8

[11] Matthias Fuchs, Bert Jüttler, Otmar Scherzer, and Huaiping Yang, *Shape metrics based on elastic deformations*, J. Math. Imaging Vis., 35 (2009), pp. 86–102. 2, 7

[12] E. S. Gawlik, P. Mullen, D. Pavlov, J. E. Marsden, and M. Desbrun, *Geometric, variational discretization of continuum theories*, Phys. D, 240 (2011), pp. 1724–1760. 2

[13] Ernst Hairer, Christian Lubich, and Gerhard Wanner, *Geometric numerical integration*, vol. 31 of Springer Series in Computational Mathematics, Springer-Verlag, Berlin, second ed., 2006. Structure-preserving algorithms for ordinary differential equations. 2

[14] B. Heeren, M. Rumpf, M. Wardetzky, and B. Wirth, *Time-discrete geodesics in the space of shells*, Computer Graphics Forum, 31 (2012), pp. 1755–1764. 8

[15] L. Kharevych, Weiwei Yang, Y. Tong, E. Kanso, J. E. Marsden, P. Schröder, and M. Desbrun, *Geometric, variational integrators for computer animation*, in Proceedings of the 2006 ACM SIGGRAPH/Eurographics symposium on Computer animation, SCA '06, Aire-la-Ville, Switzerland, Switzerland, 2006, Eurographics Association, pp. 43–51. 2

[16] Arkady Kheyfets, Warner A. Miller, and Gregory A. Newton, *Schild's ladder parallel transport procedure for an arbitrary connection*, Internat. J. Theoret. Phys., 39 (2000), pp. 2891–2898. 5

[17] M. Kilian, N. J. Mitra, and H. Pottmann, *Geometric modeling in shape space*, in ACM Transactions on Graphics, vol. 26, 2007, pp. #64, 1–8. 1, 2

[18] E. Klassen, A. Srivastava, W. Mio, and S. H. Joshi, *Analysis of planar shapes using geodesic paths on shape spaces*, IEEE Transactions on Pattern Analysis and Machine Intelligence, 26 (2004), pp. 372–383. 2

[19] A. Lew, M. Marsden, M. Oritz, and M. West, *Variational time integrators*, Int. J. Numer. Meth. Engng, 60 (2004), pp. 153–212. 2

[20] Xiuwen Liu, Yonggang Shi, Ivo Dinov, and Washington Mio, *A computational model of multidimensional shape*, International Journal of Computer Vision, Online First (2010). 2

[21] Peter W. Michor and David Mumford, *Riemannian geometries on spaces of plane curves*, J. Eur. Math. Soc., 8 (2006), pp. 1–48. 2

[22] S. Müller and M. Ortiz, *On the* Γ*-convergence of discrete dynamics and variational integrators*, J. Nonlinear Sci., 14 (2004), pp. 279–296. 2

[23] Sujit Nair, *Time adaptive variational integrators: a space-time geodesic approach*, Phys. D, 241 (2012), pp. 315–325. 2

[24] J. Nečas and M. Šilhavý, *Multipolar viscous fluids*, Quarterly of Applied Mathematics, 49 (1991), pp. 247–265. 9

[25] Sina Ober-Blöbaum, Oliver Junge, and Jerrold E. Marsden, *Discrete mechanics and optimal control: an analysis*, ESAIM Control Optim. Calc. Var., 17 (2011), pp. 322–352. 2

[26] Martin Rumpf and Benedikt Wirth, *Discrete geodesic calculus in shape space*, submitted, (2012). 1, 6

[27] F. R. Schmidt, M. Clausen, and D. Cremers, *Shape matching by variational computation of geodesics on a manifold*, in Pattern Recognition, vol. 4174 of LNCS, Springer, 2006, pp. 142–151. 2

[28] Anuj Srivastava, Aastha Jain, Shantanu Joshi, and David Kaziska, *Statistical shape models using elastic-string representations*, in Asian Conference on Computer Vision, P.J. Narayanan, ed., vol. 3851 of LNCS, 2006, pp. 612–621. 2

[29] Ganesh Sundaramoorthi, Andrea Mennucci, Stefano Soatto, and Anthony Yezzi, *A new geometric metric in the space of curves, and applications to tracking deforming objects by prediction and filtering*, SIAM Journal on Imaging Sciences, 4 (2011), pp. 109–145. 1, 2

[30] Benedikt Wirth, Leah Bar, Martin Rumpf, and Guillermo Sapiro, *A continuum mechanical approach to geodesics in shape space*, IJCV, 93 (2011), pp. 293–318. 2, 6, 7

[31] Laurent Younes, Peter W. Michor, Jayant Shah, and David Mumford, *A metric on shape space with explicit geodesics*, Atti Accad. Naz. Lincei Cl. Sci. Fis. Mat. Natur. Rend. Lincei (9) Mat. Appl., 19 (2008), pp. 25–57. 1, 2